

1.4. Computer checking of the subgroup data

BY FRANZ GÄHLER

1.4.1. Introduction

Most of the data in Part 2 of this volume have been checked by a computer-algebra program. This program is built upon the package *Cryst* (Eick *et al.*, 2001), formerly known as *CrystGAP* (Eick *et al.*, 1997), which in turn is an extension of the computer-algebra program *GAP* (The *GAP* Group, 2002). *GAP* is a program system for computations involving general algebraic structures, in particular groups. Since space groups are infinite groups, the generic algorithms for finite groups often cannot be used for space groups. The *Cryst* package provides the special methods and algorithms needed for working with space groups, thereby extending the field of applicability of *GAP*. The algorithms used in *Cryst* have been described by Eick *et al.* (1997).

1.4.2. Basic capabilities of the *Cryst* package

Before we describe in more detail the checks that have been performed, we briefly summarize the capabilities of the *Cryst* package and how space groups are handled by *Cryst*.

In *Cryst*, a space group is represented as a fixed group of augmented matrices \mathbb{W}_i , cf. Section 1.2.2.4. The group multiplication therefore coincides with matrix multiplication. The space group is defined by an arbitrary (finite) set of generating elements. The space group consists of the set of all augmented matrices that can be obtained as products of the generating matrices. It is not important which generating elements are chosen. Different sets of generators can define the same space group.

The representation of a space group as a group of augmented matrices implicitly requires the choice of an origin and a vector basis of Euclidean space. The *Cryst* package does not require any particular choice of origin or vector basis. In particular, it is not necessary to work with a lattice basis; in fact, any basis can be used. This has the advantage that one can continue to use the basis chosen for the parent group when passing to a subgroup.

Space groups are conjugated by some affine coordinate transformation if they differ only in their settings, *i.e.* by the choice of their origins and/or vector bases. Such an affine coordinate transformation is also represented by an augmented matrix. Given a coordinate transformation \mathbb{V} and a space group \mathcal{G} , the transformed space group \mathcal{G}' is simply generated by the transformed generators of \mathcal{G} .

If a space group \mathcal{G} is given by a set of generators, *Cryst* first needs to compute the point group \mathcal{P} of \mathcal{G} , and the translation subgroup \mathcal{T} , represented by a (canonical) basis of the translation lattice (the basis of the translation lattice chosen by *Cryst* is canonical in the sense that the same basis is always chosen for a given \mathcal{T}). The point group \mathcal{P} is simply generated by the linear parts of the generators of \mathcal{G} . The mapping \mathbb{H} , which sends each element of \mathcal{G} to its linear part, is in fact a group homomorphism $\mathbb{H}: \mathcal{G} \rightarrow \mathcal{P}$. The point group \mathcal{P} is finite and each point-group element p can easily be expressed as a product of generators of \mathcal{P} , which are images of generators of \mathcal{G} under \mathbb{H} . This provides a way to compute, for any $p \in \mathcal{P}$, some representative pre-image g of p under \mathbb{H} , *i.e.* some element $g \in \mathcal{G}$ which is mapped onto p by \mathbb{H} . Having expressed the element p as a product of generators, one simply replaces the

factors in this product by their representative pre-images under \mathbb{H} , which are known by construction, to obtain the representative pre-image g of p .

To compute a generating set of the translation subgroup $\mathcal{T} \leq \mathcal{G}$, one first computes a set of defining relations of the point group \mathcal{P} for the given generators. For a finite group this is a standard task in *GAP*. These defining relations are a set of inequivalent ways to express the identity element of \mathcal{P} by the generators of \mathcal{P} . Replacing the factors in such a defining relation by their representative pre-images in \mathcal{G} yields a pre-image of the identity of \mathcal{P} , *i.e.* an element of \mathcal{T} . The translations so obtained generate, together with the pure translation generators of \mathcal{G} , the entire translation subgroup $\mathcal{T} \leq \mathcal{G}$.

We now have all the necessary information to test whether a given augmented matrix \mathbb{A} is an element of a space group \mathcal{G} . One first tests whether the linear part \mathbb{M} of \mathbb{A} is an element of the point group \mathcal{P} of \mathcal{G} . If $\mathbb{M} \notin \mathcal{P}$, \mathbb{A} is not an element of \mathcal{G} . Otherwise, some pre-image \mathbb{G} of \mathbb{M} in \mathcal{G} is computed. \mathbb{A} is then an element of \mathcal{G} if and only if \mathbb{A} and \mathbb{G} differ in their translation part by an element of the translation group \mathcal{T} . With this membership test, we can determine whether two space groups are equal, or whether one is a subgroup of the other: a space group \mathcal{H} is a subgroup of a space group \mathcal{G} if all the generators of \mathcal{H} are elements of \mathcal{G} . Two space groups are equal if either of them is a subgroup of index 1 of the other.

1.4.3. Computing maximal subgroups

The *Cryst* package has built-in facilities for computing the maximal subgroups of a given index for any space group \mathcal{G} . More precisely, given a prime number p , *Cryst* can compute conjugacy-class representatives of those maximal subgroups of \mathcal{G} whose index in \mathcal{G} is a power of p . The algorithms used for this task are described in Eick *et al.* (1997). Essentially, one determines the maximal subgroups of the (finite) factor group $\mathcal{G}/\mathcal{T}_p$, where \mathcal{T}_p is the subgroup of those translations of \mathcal{G} which are a p -fold multiple of an element of the full translation group \mathcal{T} of \mathcal{G} . After the maximal subgroups are obtained, the translations in \mathcal{T}_p are added back to the subgroups.

From a representative \mathcal{H} of a conjugacy class of subgroups, the list of all subgroups in the same conjugacy class is obtained by repeatedly conjugating \mathcal{H} (and the subgroups obtained from it by conjugation) with generators of \mathcal{G} , until no new conjugated subgroups are obtained. This is also the way of determining whether two subgroups are conjugated: one enumerates the groups in the conjugacy class of one of them, and checks whether the other is among them.

The index of a subgroup \mathcal{H} of \mathcal{G} is easily computed as the product of the index of the point group of \mathcal{H} in the point group of \mathcal{G} and the index of the translation group of \mathcal{H} in the translation group of \mathcal{G} . For maximal subgroups, only one of these factors is different from 1. For *klassenleiche* subgroups, the two point groups are equal, whereas for *translationenleiche* subgroups the two translation subgroups are equal. Therefore, a maximal subgroup is easily identified either as a *klassenleiche* or a *translationenleiche* subgroup.

1. SPACE GROUPS AND THEIR SUBGROUPS

1.4.4. Description of the checks

In order to avoid new errors being introduced in the typesetting process after the data have been checked and corrected, we carried out the computer checks directly on the $\text{\LaTeX} 2_{\epsilon}$ sources used for the production of this volume. The tables have been typeset with specially designed $\text{\LaTeX} 2_{\epsilon}$ macros, the primary purpose of which was to guarantee a homogeneous layout throughout the book. As a side effect, it was relatively easy to write a *GAP* program which parses the $\text{\LaTeX} 2_{\epsilon}$ sources, recognizes the macros, extracts the data (the arguments of the macros), brings the data into a form suitable for further processing with routines from the *Cryst* package and finally performs the various checks. In this way, the checks could be done fully automatically, and could be repeated after every modification of the tabulated data.

In the following, we describe the checks that have been carried out. We first applied a number of tests individually to each of the tabulated maximal subgroups of low index: whether it is a subgroup, whether the index given is correct, whether the listed coordinate transformation maps the subgroup to the preferred setting of the given space-group type (and thus, whether the space-group

type of the subgroup is correct) and whether the listed coordinate transformation maps the given generators of the subgroup to the standard generators of its space-group type, in the same order. Here, the preferred setting is the setting of the parent group, where applicable, and otherwise the preferred setting of the space-group type of the subgroup, if there is more than one setting in the tables.

In a second step, a complete set of maximal subgroups of low index (2, 3 or 4) is computed afresh with the routines from the *Cryst* package. These subgroups are then divided into conjugacy classes, and classified as *klassenleiche* or *translationenleiche* subgroups. This list is then compared with the tabulated list of maximal subgroups. It was verified that each maximal subgroup of a given index was listed exactly once, that the classification into conjugacy classes of subgroups was correct and that the subgroups were correctly identified as *klassenleiche* or as *translationenleiche* subgroups.

All the tests described above concern the maximal subgroups of low index. Unfortunately, similar automatic tests could not be performed on the series of isomorphic subgroups. The subgroups in these series contain variable parameters, and *Cryst* can only deal with fixed, concrete space groups without free parameters.

References

- Aroyo, M. I. & Perez-Mato, J. M. (1998). *Symmetry-mode analysis of displacive phase transitions using International Tables for Crystallography*. *Acta Cryst.* **A54**, 19–30.
- Ascher, E. (1966). *Role of particular maximal subgroups in continuous phase transitions*. *Phys. Lett.* **20**, 352–354.
- Ascher, E. (1967). *Symmetry changes in continuous transitions. A simplified theory applied to V_3Si* . *Chem. Phys. Lett.* **1**, 69–72.
- Ascher, E. (1968). *Lattices of equi-translation subgroups of the space groups*. Geneva: Battelle Institute.
- Ascher, E., Gramlich, V. & Wondratschek, H. (1969). *Corrections to the sections 'Untergruppen' of the space groups in Internationale Tabellen zur Bestimmung von Kristallstrukturen (1935) Vol. I*. *Acta Cryst.* **B25**, 2154–2156.
- Astbury, W. T. & Yardley, K. (1924). *Tabulated data for the examination of the 230 space-groups by homogeneous X-rays*. *Philos. Trans. R. Soc. London*, **224**, 221–257.
- Barlow, W. (1894). *Über die geometrischen Eigenschaften homogener starrer Strukturen*. *Z. Kristallogr. Mineral.* **23**, 1–63.
- Bärnighausen, H. (1975). *Group-subgroup relations between space groups as an ordering principle in crystal chemistry: the "family tree" of perovskite-like structures*. *Acta Cryst.* **A31**, part S3, S3, 01.1-9.
- Bärnighausen, H. (1980). *Group-subgroup relations between space groups: a useful tool in crystal chemistry*. *MATCH Comm. Math. Chem.* **9**, 139–175.
- Baur, W. H. (1994). *Rutile type derivatives*. *Z. Kristallogr.* **209**, 143–150.
- Baur, W. H. & Kassner, D. (1992). *The perils of Cc: comparing the frequencies of falsely assigned space groups with their general population*. *Acta Cryst.* **B48**, 356–369.
- Baur, W. H. & McLarnan, T. J. (1982). *Observed wurtzite derivatives and related tetrahedral structures*. *J. Solid State Chem.* **42**, 300–321.
- Bertaut, E. F. & Billiet, Y. (1979). *On equivalent subgroups and supergroups of the space groups*. *Acta Cryst.* **A35**, 733–745.
- Billiet, Y. (1973). *Les sous-groupes isosymboliques des groupes spatiaux*. *Bull. Soc. Fr. Minéral. Cristallogr.* **96**, 327–334.
- Billiet, Y. (1980). *The subgroups of finite index of the space groups: determination via conventional coordinate systems*. *MATCH Comm. Math. Chem.* **9**, 127–190.
- Billiet, Y. & Bertaut, E. F. (2002). *Isomorphic subgroups of space groups*. *International Tables for Crystallography*, Vol. A, *Space-group symmetry*, edited by Th. Hahn, Part 13. Dordrecht: Kluwer Academic Publishers.
- Billiet, Y. & Rolley Le Coz, M. (1980). *Le groupe $P1$ et ses sous-groupes. II. Tables des sous-groupes*. *Acta Cryst.* **A36**, 242–248.
- Billiet, Y. & Sayari, A. (1984). *Les sous-groupes isomorphes d'un group d'espace de type $p4$. I. Détermination univoque*. *Acta Cryst.* **A40**, 624–631.
- Billiet, Y., Sayari, A. & Zarrouk, H. (1978). *L'association des familles de Wyckoff dans les changements de repère conventionnel des groupes spatiaux et les passages aux sous-groupes spatiaux*. *Acta Cryst.* **A34**, 811–819.
- Biltz, W. (1934). *Raumchemie der festen Stoffe*. Leipzig: L. Voss.
- Birman, J. L. (1966a). *Full group and subgroup methods in crystal physics*. *Phys. Rev.* **150**, 771–782.
- Birman, J. L. (1966b). *Simplified theory of symmetry change in second-order phase transitions: application to V_3Si* . *Phys. Rev. Lett.* **17**, 1216–1219.
- Bock, O. & Müller, U. (2002a). *Symmetrieverwandtschaften bei Varianten des Perowskit-Typs*. *Acta Cryst.* **B58**, 594–606.
- Bock, O. & Müller, U. (2002b). *Symmetrieverwandtschaften bei Varianten des ReO_3 -Typs*. *Z. Anorg. Allg. Chem.* **628**, 987–992.
- Boyle, L. L. & Lawrenson, J. E. (1973). *The origin dependence of the Wyckoff site description of a crystal structure*. *Acta Cryst.* **A29**, 353–357.
- Bravais, A. (1850). *Mémoire sur les systèmes formés par les points distribués régulièrement sur un plan ou dans l'espace*. *J. Ecole Polytech.* **19**, 1–128. (English: Memoir 1, Crystallographic Society of America, 1949.)
- Brown, H., Bülow, R., Neubüser, J., Wondratschek, H. & Zassenhaus, H. (1978). *Crystallographic groups of four-dimensional space*. New York: John Wiley & Sons.
- Buerger, M. J. (1947). *Derivative crystal structures*. *J. Chem. Phys.* **15**, 1–16.
- Buerger, M. J. (1951). *Phase transformations in solids*, ch. 6. New York: Wiley.
- Burckhardt, J. J. (1988). *Die Symmetrie der Kristalle*. Basel: Birkhäuser.
- Byström, A., Hök, B. & Mason, B. (1941). *The crystal structure of zinc metaantimonate and similar compounds*. *Ark. Kemi Mineral. Geol.* **15B**, 1–8.
- Chapuis, G. C. (1992). *Symmetry relationships between crystal structures and their practical applications. Modern perspectives in inorganic chemistry*, edited by E. Parthé, pp. 1–16. Dordrecht: Kluwer Academic Publishers.
- Cracknell, A. P. (1975). *Group theory in solid state physics*. New York: Taylor and Francis Ltd. and Pergamon.
- Deonarine, S. & Birman, J. L. (1983). *Group-subgroup phase transitions, Hermann's space group decomposition theorem, and chain subduction criterion in crystals*. *Phys. Rev. B*, **27**, 4261–4273.
- Eick, B., Gähler, F. & Nickel, W. (1997). *Computing maximal subgroups and Wyckoff positions of space groups*. *Acta Cryst.* **A53**, 467–474.
- Eick, B., Gähler, F. & Nickel, W. (2001). *The 'Cryst' Package*. Version 4.1. <http://www.itap.physik.uni-stuttgart.de/~gaehler/gap/packages.html>.
- Fedorov, E. (1891). *Symmetry of regular systems and figures*. *Zap. Mineral. Obshch.* (2), **28**, 1–46. (In Russian.) (English: *Symmetry of crystals*, American Crystallographic Association, 1971.)
- Finken, H., Neubüser, J. & Plesken, W. (1980). *Space groups and groups of prime power order II*. *Arch. Math.* **35**, 203–209.
- Fischer, W. & Koch, E. (1983). *On the equivalence of point configurations due to Euclidean normalizers (Cheshire groups) of space groups*. *Acta Cryst.* **A39**, 907–915.
- Flack, H. D., Wondratschek, H., Hahn, Th. & Abrahams, S. C. (2000). *Symmetry elements in space groups and point groups. Addenda to two IUCr reports on the nomenclature of symmetry*. *Acta Cryst.* **A56**, 96–98.
- Goldschmidt, V. M. (1926). *Untersuchungen über den Bau und Eigenschaften von Kristallen*. *Skr. Nor. Vidensk. Akad. Oslo Mat.-Nat. Kl.* 1926 No. 2 and 1927 No. 8.
- Haag, F. (1887). *Die regulären Kristallkörper. Programm des Königlichen Gymnasiums in Rottweil zum Schlusse des Schuljahres 1886–1887*.
- Hahn, Th. & Wondratschek, H. (1994). *Symmetry of crystals. Introduction to International Tables for Crystallography*, Vol. A. Sofia: Heron Press.
- Heesch, H. (1930). *Zur systematischen Strukturtheorie. II*. *Z. Kristallogr.* **72**, 177–201.
- Hermann, C. (1929). *Zur systematischen Strukturtheorie. IV. Untergruppen*. *Z. Kristallogr.* **69**, 533–555.
- International Tables for Crystallography* (1983). Vol. A, *Space-group symmetry*, edited by Th. Hahn, 1st ed. Dordrecht: Kluwer Academic Publishers. (Abbreviated *IT A*.)
- International Tables for Crystallography* (2002). Vol. A, *Space-group symmetry*, edited by Th. Hahn, 5th ed. Dordrecht: Kluwer Academic Publishers. (Abbreviated *IT A*.)
- International Tables for Crystallography* (2002). Vol. E, *Subperiodic groups*, edited by V. Kopský & D. B. Litvin. Dordrecht: Kluwer Academic Publishers.
- International Tables for X-ray Crystallography* (1952, 1965, 1969). Vol. I, *Symmetry groups*, edited by N. F. M. Henry & K. Lonsdale. Birmingham: Kynoch Press.
- Internationale Tabellen zur Bestimmung von Kristallstrukturen* (1935). 1. Bd. Edited by C. Hermann. Berlin: Borntraeger. (In German, English and French.) (Abbreviated *IT 35*.)
- Izumov, Y. A. & Syromyatnikov, V. N. (1990). *Phase transitions and crystal symmetry*. Dordrecht: Kluwer Academic Publishers.
- Janovec, V. (1976). *A symmetry approach to domain structures*. *Ferroelectrics*, **12**, 43–53.

REFERENCES

- Janovec, V. & Přívratská, J. (2003). *Domain structures. International tables for crystallography*, Vol. D, edited by A. Authier, *Physical properties of crystals*, ch. 3.4. Dordrecht: Kluwer Academic Publishers.
- Koch, E. (1984). *The implications of normalizers on group-subgroup relations between space groups*. *Acta Cryst.* **A40**, 593–600.
- Koch, E. & Fischer, W. (1975). *Automorphismengruppen von Raumgruppen und die Zuordnung von Punktlagen zu Konfigurationslagen*. *Acta Cryst.* **A31**, 88–95.
- Koch, E. & Fischer, W. (1985). *Lattice complexes and limiting complexes versus orbit types and non-characteristic orbits: a comparative discussion*. *Acta Cryst.* **A41**, 421–426.
- Koch, E., Fischer, W. & Müller, U. (2002). *Normalizers of space groups and their use in crystallography. International tables for crystallography*, Vol. A, *Space-group symmetry*, edited by Th. Hahn, Part 15. Dordrecht: Kluwer Academic Publishers.
- Kroumova, E., Aroyo, M. I. & Pérez-Mato, J. M. (2002). *Prediction of new displacive ferroelectrics through systematic pseudosymmetry search. Results for materials with Pba2 and Pmc2₁ symmetry*. *Acta Cryst.* **B58**, 921–933.
- Kroumova, E., Perez-Mato, J. M. & Aroyo, M. I. (1998). *WYCKSPLIT. Computer program for the determination of the relations of Wyckoff positions for a group-subgroup pair*. *J. Appl. Cryst.* **31**, 646. Accessible at: <http://www.cryst.ehu.es/cryst/wpsplit.html>.
- Lampart, L. (1994). *A document preparation system*. 2nd ed. Reading, MA: Addison-Wesley.
- Landau, L. D. (1937). *Theory of phase transitions*. *Zh. Eksp. Teoret. Fiz.* **7**, pp. 19–32, 627–636. (In Russian.) (German: *Phys. Z. Sowjetunion*, **11**, pp. 20–47, 545–555.)
- Landau, L. D. & Lifshitz, E. M. (1980). *Statistical physics*, Part 1, 3rd ed., pp. 459–471. London: Pergamon. (Russian: *Statisticheskaya Fizika*, chast 1. Moskva: Nauka, 1976; German: *Lehrbuch der theoretischen Physik*, 6. Aufl., Bd. 5, Teil 1, S. 436–447. Berlin: Akademie-Verlag, 1984.)
- Lawrenson, J. E. (1972). *Theoretical studies in crystallography*. Dissertation, University of Kent at Canterbury.
- Ledermann, W. (1976). *Introduction to group theory*. London: Longman. (German: *Einführung in die Gruppentheorie*, Braunschweig: Vieweg, 1977.)
- Levanyuk, A. P. & Sannikov, D. G. (1971). *Phenomenological theory of dielectric anomalies in ferroelectric materials with several phase transitions at temperatures close together*. *Sov. Phys. JETP*, **11**, 600–604.
- McLarnan, T. J. (1981a). *Mathematical tools for counting polytypes*. *Z. Kristallogr.* **155**, 227–245.
- McLarnan, T. J. (1981b). *The numbers of polytypes in close packings and related structures*. *Z. Kristallogr.* **155**, 269–291.
- McLarnan, T. J. (1981c). *The combinatorics of cation-deficient close-packed structures*. *J. Solid State Chem.* **26**, 235–244.
- Marsh, R. E., Kapon, M., Hu, S. & Herbstein, F. H. (2002). *Some 60 new space-group corrections*. *Acta Cryst.* **B58**, 62–77.
- Megaw, H. D. (1973). *Crystal structures: A working approach*. Philadelphia: Saunders.
- Meyer, A. (1981). *Symmetriebeziehungen zwischen Kristallstrukturen des Formeltyps AX₂, ABX₄ und AB₂X₆ sowie deren Ordnungs- und Leerstellenvarianten*. Dissertation, Universität Karlsruhe.
- Müller, U. (1978). *Strukturmöglichkeiten für Pentahalogenide mit Doppeloktaeder-Molekülen (MX₅)₂ bei dichtester Packung der Halogenatome*. *Acta Cryst.* **A34**, 256–267.
- Müller, U. (1980). *Strukturverwandtschaften unter den EPh₄⁺-Salzen*. *Acta Cryst.* **B36**, 1075–1081.
- Müller, U. (1981). *MX₄-Ketten aus kantenverknüpften Oktaedern: mögliche Kettenkonfigurationen und mögliche Kristallstrukturen*. *Acta Cryst.* **B37**, 532–545.
- Müller, U. (1986). *MX₅-Ketten aus eckenverknüpften Oktaedern. Mögliche Kettenkonfigurationen und mögliche Kristallstrukturen bei dichtester Packung der X-Atome*. *Acta Cryst.* **B42**, 557–564.
- Müller, U. (1992). *Berechnung der Anzahl möglicher Strukturtypen für Verbindungen mit dichtest gepackter Anionenteilstruktur. I. Das Rechenverfahren*. *Acta Cryst.* **B48**, 172–178.
- Müller, U. (1993). *Inorganic structural chemistry*, pp. 233–246. Chichester, New York: Wiley. (German: *Anorganische Strukturchemie*, 3. Aufl., 1996, S. 296–309. Stuttgart: Teubner.)
- Müller, U. (1998). *Strukturverwandtschaften zwischen trigonalen Verbindungen mit hexagonal-dichtester Anionenteilstruktur und besetzten Oktaederlücken*. *Z. Anorg. Allg. Chem.* **624**, 529–532.
- Müller, U. (2002). *Kristallpackungen mit linear koordinierten Atomen*. *Z. Anorg. Allg. Chem.* **628**, 1269–1278.
- Müller, U. (2003). *Die Zahl der Substitutions- und Leerstellenvarianten des NaCl-Typs bei verdoppelter Elementarzelle (a, b, 2c)*. *Z. Anorg. Allg. Chem.* **629**, 487–492.
- Müller, U. & Brelle, A. (1995). *Über isomorphe Untergruppen von Raumgruppen der Kristallklassen 4, $\bar{4}$, 4/m, 3, $\bar{3}$, 6, $\bar{6}$ und 6/m*. *Acta Cryst.* **A51**, 300–304.
- Nespolo, M. & Ferraris, G. (2004). *Applied geminography – symmetry analysis of twinned crystals and definition of twinning by reticular polyhohedry*. *Acta Cryst.* **A60**, 89–95.
- Neubüser, J. & Wondratschek, H. (1966). *Untergruppen der Raumgruppen*. *Krist. Tech.* **1**, 529–543.
- Niggli, P. (1919). *Geometrische Kristallographie des Diskontinuums*. Leipzig: Gebrüder Borntraeger. Reprint (1973) Wiesbaden: Dr M. Saendig.
- Nolze, G. (1996). *POWDER CELL. Computer program for the calculation of X-ray powder diagrams*. Bundesanstalt für Materialforschung, Berlin.
- Pauling, L. (1928). *The coordination theory of the structure of ionic compounds – Probleme der modernen Physik (Sommerfeld-Festschrift)*, p. 11. Leipzig: S. Hirzel.
- Pauling, L. (1929). *The principles determining the structures of complex ionic crystals*. *J. Am. Chem. Soc.* **51**, 1010–1026.
- Plesken, W. & Schulz, T. (2000). *Counting crystallographic groups in low dimensions*. *Exp. Math.* **9**, 407–411.
- Pöttgen, R. & Hoffmann, R.-D. (2001). *AlB₂-related intermetallic compounds – a comprehensive view based on a group-subgroup scheme*. *Z. Kristallogr.* **216**, 127–145.
- Salje, E. K. H. (1990). *Phase transitions in ferroelastic and co-elastic crystals*. Cambridge University Press.
- Sayari, A. (1976). *Contribution à l'étude des groupes d'espace bidimensionnels, dérivation des sous-groupes et des surstructures*. Thesis, University of Tunis, Tunisia.
- Sayari, A. & Billiet, Y. (1977). *Subgroups and changes of standard settings of triclinic and monoclinic space groups*. *Acta Cryst.* **A33**, 985–986.
- Schoenflies, A. M. (1891). *Krystallsysteme und Krystallstruktur*. Leipzig: Teubner. Reprint 1984: Springer.
- Sohncke, L. (1879). *Entwicklung einer Theorie der Krystallstruktur*. Leipzig: Teubner.
- Sowa, H. (2001). *On the transition from the wurtzite to the NaCl type*. *Acta Cryst.* **A57**, 176–182.
- Stokes, H. T. & Hatch, D. M. (1988). *Isotropy subgroups of the 230 crystallographic space groups*. Singapore: World Scientific.
- Stokes, H. T. & Hatch, D. M. (2002). *Procedure for obtaining microscopic mechanisms of reconstructive phase transitions in crystalline solids*. *Phys. Rev. B*, **65**, 144114–144123.
- Strukturbericht 1913–1928* (1931). Edited by P. P. Ewald & C. Hermann. Leipzig: Akademische Verlagsgesellschaft. Continued by *Structure Reports*.
- Tendeloo, G. van & Amelinckx, S. (1974). *Group-theoretical considerations concerning domain formation in ordered alloys*. *Acta Cryst.* **A30**, 431–440.
- The GAP Group (2002). *GAP – Groups, Algorithms and Programming*. Version 4.3. <http://www.gap-system.org>.
- Tolédano, J.-C., Janovec, V., Kopský, V., Scott, J.-F. & Boček, P. (2003). *Structural phase transitions. International tables for crystallography*, Vol. D, *Physical properties of crystals*, edited by A. Authier, ch. 3.1. Dordrecht: Kluwer Academic Publishers.
- Tolédano, J.-C. & Tolédano, P. (1987). *The Landau theory of phase transitions*. Singapore: World Scientific.
- Wiener, C. (1863). *Grundzüge der Weltordnung. I. Atomlehre*, p. 82. Leipzig: Heidelberg.
- Wondratschek, H. (1976). *Extraordinary orbits of space groups. Theoretical considerations*. *Z. Kristallogr.* **143**, 460–470.
- Wondratschek, H. (1980). *Crystallographic orbits, lattice complexes, and orbit types*. *MATCH Comm. Math. Chem.* **9**, 121–125.

REFERENCES

- Wondratschek, H. (1993). *Splitting of Wyckoff positions (orbits)*. *Mineral. Petrol.* **48**, 87–96.
- Wondratschek, H. (2002). *Special topics on space groups*. *International tables for crystallography*, Vol. A, *Space-group symmetry*, edited by Th. Hahn, Part 8. Dordrecht: Kluwer Academic Publishers.
- Wondratschek, H. & Aroyo, M. I. (2001). *The application of Hermann's group \mathcal{M} in group–subgroup relations between space groups*. *Acta Cryst.* **A57**, 311–320.
- Wondratschek, H. & Jeitschko, W. (1976). *Twin domains and antiphase domains*. *Acta Cryst.* **A32**, 664–666.
- Wondratschek, H., Müller, U., Aroyo, M. I. & Sens, I. (1995). *Splitting of Wyckoff positions (orbits)*. II. *Group–subgroup chains of index 6*. *Z. Kristallogr.* **210**, 567–573.
- Wyckoff, R. W. G. (1922). *The analytical expression of the results of the theory of space groups*. Washington: Carnegie Institution.
- Wyckoff, R. W. G. (1965). *Crystal structures*, 2nd ed., Vol. 3, pp. 361–362. New York: Interscience.