

2. MAXIMAL SUBGROUPS OF THE PLANE GROUPS AND SPACE GROUPS

empty subblock is then designated by 'none'; in the other subblock the supergroups are listed. The kind of listing depends on the subblock. Examples may be found in the tables of $P222$, No. 16, and $Fd\bar{3}c$, No. 228.

Under the heading 'Additional centring translations', the supergroups are listed by their indices and either by their nonconventional HM symbols, with the space-group numbers and the standard HM symbols in parentheses, or by their conventional HM symbols and only their space-group numbers in parentheses. Examples are provided by space group $Pbca$, No. 61, with both subblocks non-empty and by space group $P222$, No. 16, with supergroups only under the heading 'Additional centring translations'.

Under the heading 'Decreased unit cell' each supergroup is listed by its index and by its lattice relations, where the basis vectors \mathbf{a}' , \mathbf{b}' and \mathbf{c}' refer to the supergroup \mathcal{G} and the basis vectors \mathbf{a} , \mathbf{b} and \mathbf{c} to the original group \mathcal{H} . After these data are listed either the nonconventional HM symbol, followed by the space-group number and the conventional HM symbol in parentheses, or the conventional HM symbol with the space-group number in parentheses. Examples are provided again by space group $Pbca$, No. 61, with both subblocks occupied and space group $F\bar{4}3m$, No. 216, with an empty subblock 'Additional centring translations' but data under the heading 'Decreased unit cell'.

2.1.6.4. Isomorphic supergroups

Each space group \mathcal{G} has an infinite number of isomorphic subgroups \mathcal{H} because the number of primes is infinite. For the same reason, each space group \mathcal{H} has an infinite number of isomorphic supergroups \mathcal{G} . They are not listed in the tables of this volume because they are implicitly listed among the subgroup data.

2.1.7. The subgroup graphs

2.1.7.1. General remarks

The group–subgroup relations between the space groups may also be described by graphs. This way is chosen in Chapters 2.4 and 2.5. Graphs for the group–subgroup relations between crystallographic point groups have been published, for example, in *Internationale Tabellen zur Bestimmung von Kristallstrukturen* (1935) and in *IT A* (2002), Fig. 10.1.4.3. Three kinds of graphs for subgroups of space groups have been constructed and can be found in the literature:

- (1) Graphs for t -subgroups, such as the graphs of Ascher (1968).
- (2) Graphs for k -subgroups, such as the graphs for cubic space groups of Neubüser & Wondratschek (1966).
- (3) Mixed graphs, combining t - and k -subgroups. These are used, for example, when relations between existing or suspected crystal structures are to be displayed. An example is the 'family tree' of Bärnighausen (1980), Fig. 15, now called a *Bärnighausen tree*.

A complete collection of graphs of the first two kinds is presented in this volume: in Chapter 2.4 those displaying the *translationengleiche* or t -subgroup relations and in Chapter 2.5 those for the *klassengleiche* or k -subgroup relations. Neither type of graph is restricted to maximal subgroups but both contain t - or k -subgroups of higher indices, with the exception of isomorphic subgroups, cf. Section 2.1.7.3 below.

The group–subgroup relations are direct relations between the space groups themselves, not between their types. However, each such relation is valid for a pair of space groups, one from each of

the types, and for each space group of a given type there exists a corresponding relation. In this sense, one can speak of a relation between the space-group types, keeping in mind the difference between space groups and space-group types, cf. Section 1.2.5.3.

The space groups in the graphs are denoted by the standard HM symbols and the space-group numbers. In each graph, each space-group type is displayed at most once. Such graphs are called *contracted graphs* here. Without this contraction, the more complex graphs would be much too large for the page size of this volume.

The symbol of a space group \mathcal{G} is connected by uninterrupted straight lines with the symbols of all maximal non-isomorphic subgroups \mathcal{H} or minimal non-isomorphic supergroups \mathcal{S} of \mathcal{G} . In general, the *maximal subgroups* of \mathcal{G} are drawn on a *lower level* than \mathcal{G} ; in the same way, the *minimal supergroups* of \mathcal{G} are mostly drawn on a *higher level* than \mathcal{G} . For exceptions see Section 2.1.7.3. Multiple lines may occur in the graphs for t -subgroups. They are explained in Section 2.1.7.2. No indices are attached to the lines. They can be taken from the corresponding subgroup tables of Chapter 2.3, and are also provided by the general formulae of Section 1.2.8. For the k -subgroup graphs, they are further specified at the end of Section 2.1.7.3.

2.1.7.2. Graphs for translationengleiche subgroups

Let \mathcal{G} be a space group and $\mathcal{T}(\mathcal{G})$ the normal subgroup of all its translations. Owing to the isomorphism between the factor group $\mathcal{G}/\mathcal{T}(\mathcal{G})$ and the point group $\mathcal{P}_{\mathcal{G}}$, see Section 1.2.5.4, according to the first isomorphism theorem, Ledermann (1976), t -subgroup graphs are the same (up to the symbols) as the corresponding graphs between point groups. However, in this volume, the graphs are not complete but are contracted by displaying each space-group type at most once. This contraction may cause the graphs to look different from the point-group graphs and also different for different space groups of the same point group, cf. Example 2.1.7.2.1.

One can indicate the connections between a space group \mathcal{G} and its maximal subgroups in different ways. In the contracted t -subgroup graphs one line is drawn for each conjugacy class of maximal subgroups of \mathcal{G} . Thus, a line represents the connection to an individual subgroup only if this is a normal maximal subgroup of \mathcal{G} , otherwise it represents the connection to more than one subgroup. The conjugacy relations are not necessarily transferable to non-maximal subgroups, cf. Example 2.1.7.2.2. On the other hand, multiple lines are possible, see the examples. Although it is not in general possible to reconstruct the complete graph from the contracted one, the content of information of such a graph is higher than that of a graph which is drawn with simple lines only.

The graph for the space group at its top also contains the contracted graphs for all subgroups which occur in it, see the remark below Example 2.1.7.2.2.

Owing to lack of space for the large graphs, in all graphs of t -subgroups the group $P1$, No. 1, and its connections have been omitted. Therefore, to obtain the full graph one has to supplement the graphs by $P1$ at the bottom and to connect $P1$ by one line to each of the symbols that have no connection downwards.

Within the same graph, symbols on the same level indicate subgroups of the same index relative to the group at the top. The distance between the levels indicates the size of the index. For a more detailed discussion, see Example 2.1.7.2.2. For the sequence and the numbers of the graphs, see the paragraph below Example 2.1.7.2.2.

Example 2.1.7.2.1.

Compare the t -subgroup graphs in Figs. 2.4.4.2, 2.4.4.3 and 2.4.4.8 of $Pnna$, No. 52, $Pmna$, No. 53, and $Cmce$, No. 64, respectively. The *complete* (uncontracted) *graphs* would have the shape of the graph of the point group mmm with mmm at the top (first level), seven point groups⁵ (222 , $mm2$, $m2m$, $2mm$, $112/m$, $12/m1$ and $2/m11$) in the second level, seven point groups (112 , 121 , 211 , $11m$, $1m1$, $m11$ and $\bar{1}$) in the third level and the point group 1 at the bottom (fourth level). The group mmm is connected to each of the seven subgroups at the second level by one line. Each of the groups of the second level is connected with three groups of the third level by one line. All seven groups of the third level are connected by one line each with the point group 1 at the bottom.

The *contracted graph* of the point group mmm would have mmm at the top, three point-group types (222 , $mm2$ and $2/m$) at the second level and three point-group types (2 , m and $\bar{1}$) at the third level. The point group 1 at the bottom would not be displayed (no fourth level). Single lines would connect mmm with 222 , $mm2$ with 2 , $2/m$ with 2 , $2/m$ with m and $2/m$ with $\bar{1}$; a double line would connect $mm2$ with m ; triple lines would connect mmm with $mm2$, mmm with $2/m$ and 222 with 2 .

The number of fields in a contracted t -subgroup graph is between the numbers of fields in the full and in the contracted point-group graphs. The graph in Fig. 2.4.4.2 of $Pnna$, No. 52, has six space-group types at the second level and four space-group types at the third level. For the graph in Fig. 2.4.4.3 of $Pmna$, No. 53, these numbers are seven and five and for the graph in Fig. 2.4.4.8 of $Cmce$, No. 64 (formerly $Cmca$), the numbers are seven and six. However, in all these graphs the number of connections is always seven from top to the second level and three from each field of the second level downwards to the ground level, independent of the amount of contraction and of the local multiplicity of lines.

Example 2.1.7.2.2.

Compare the t -subgroup graphs shown in Fig. 2.4.1.1 for $Pm\bar{3}m$, No. 221, and Fig. 2.4.1.5, $Fm\bar{3}m$, No. 225. These graphs are contracted from the point-group graph $m\bar{3}m$. There are altogether nine levels (without the lowest level of $P1$). The indices relative to the top space groups $Pm\bar{3}m$ and $Fm\bar{3}m$ are 1, 2, 3, 4, 6, 8, 12, 16 and 24, corresponding to the point-group orders 48, 24, 16, 12, 8, 6, 4, 3 and 2, respectively. The height of the levels in the graphs reflects the index; the distances between the levels are slightly distorted in order to adapt to the density of the lines. From the top space-group symbol there are five lines to the symbols of maximal subgroups: The three symbols at the level of index 2 are those of cubic normal subgroups, the one (tetragonal) symbol at the level of index 3 represents a conjugacy class of three, the symbol $R\bar{3}m$, No. 166, at the level of index 4 represents a conjugacy class of four subgroups.

The graphs differ in the levels of the indices 12 and 24 (orthorhombic, monoclinic and triclinic subgroups) by the number of symbols (nine and seven for index 12, five and three for index 24). The number of lines between neighbouring connected levels depends only on the number and kind of symbols in the upper level. This property makes such graphs particularly useful.

However, for non-maximal subgroups the conjugacy relations may not hold. For example, in Fig. 2.4.1.1, the space group

$P222$ has three normal maximal subgroups of type $P2$ and is thus connected to its symbol by a triple line, although these subgroups are conjugate subgroups of the non-minimal supergroup $Pm\bar{3}m$.

The t -subgroup graphs in Figs. 2.4.1.1 and 2.4.1.5 contain the t -subgroups of $Pm\bar{3}m$ (221) and $Fm\bar{3}m$ (225) and their relations. In addition, the t -subgroup graph of $Pm\bar{3}m$ includes the t -subgroup graphs of $P432$, $P\bar{4}3m$, $Pm\bar{3}$, $P23$, $P4/mmm$, $P\bar{4}2m$, $P\bar{4}m2$, $P4mm$, $R\bar{3}m$, $R3m$ etc., that of $Fm\bar{3}m$ includes those of $F432$, $F\bar{4}3m$, $Fm\bar{3}$, $I4/mmm$, also $R\bar{3}m$ etc. Thus, many other graphs can be extracted from the two basic graphs. The same holds for the other graphs displayed in Figs. 2.4.1.2 to 2.4.4.8: each of them includes the contracted graphs of all its subgroups. For this reason one does not need 229 or 218 different graphs to cover all t -subgroup graphs of the 229 space-group types but only 37 ($P1$ can be excluded as trivial).

The preceding Example 2.1.7.2.2 suggests that one should choose the graphs in such a way that their number can be kept small. It is natural to display the ‘big’ graphs first and later those smaller graphs that are still missing. This procedure is behind the sequence of the t -subgroup graphs in this volume.

- (1) The ten graphs of $Pm\bar{3}m$, No. 221, to $Ia\bar{3}d$, No. 230, form the first set of graphs in Figs. 2.4.1.1 to 2.4.1.10.
- (2) There are a few cubic space groups left which do not appear in the first set. They are covered by the graphs of $P4_132$ (213), $P4_332$ (212) and $Pa\bar{3}$ (205). These graphs have large parts in common so that they can be united in Fig. 2.4.1.11.
- (3) No cubic space group is left now, but only eight tetragonal space groups of crystal class $4/mmm$ have appeared up to now. Among them are all graphs for $4/mmm$ space groups with an I lattice which are contained in Figs. 2.4.1.5 to 2.4.1.8 of the F -centred cubic space groups. The next 12 graphs, Figs. 2.4.2.1 to 2.4.2.12, are those for the space groups of the crystal class $4/mmm$ with lattice symbol P and different third and fourth constituents of the HM symbol. They start with $P4/mcc$, No. 124, and end with $P4_2/ncm$, No. 138.
- (4) Two (enantiomorphic) tetragonal space-group types are left which are compiled in Fig. 2.4.2.13.
- (5) The next set is formed by the four graphs in Figs. 2.4.3.1 to 2.4.3.4 of the hexagonal space groups $P6/mmm$, No. 191, to $P6_3/mmc$, No. 194. The hexagonal and trigonal enantiomorphic space groups do not appear in these graphs. They are combined in Fig. 2.4.3.5, the last one of hexagonal origin.
- (6) Several orthorhombic space groups are still left. They are treated in the eight graphs in Figs. 2.4.4.1 to 2.4.4.8, from $Pmma$, No. 51, to $Cmce$, No. 64 (formerly $Cmca$).
- (7) For each space group, the contracted graph of all its t -subgroups is provided in at least one of these 37 graphs.

For the index of a maximal t -subgroup, Lemma 1.2.8.2.3 is repeated: the index of a maximal non-isomorphic subgroup \mathcal{H} is always 2 for oblique, rectangular and square plane groups and for triclinic, monoclinic, orthorhombic and tetragonal space groups \mathcal{G} . The index is 2 or 3 for hexagonal plane groups and for trigonal and hexagonal space groups \mathcal{G} . The index is 2, 3 or 4 for cubic space groups \mathcal{G} .

2.1.7.3. Graphs for klassengleiche subgroups

There are 29 graphs for *klassengleiche* or k -subgroups, one for each crystal class with the exception of the crystal classes 1 , $\bar{1}$ and $\bar{6}$ with only one space-group type each: $P1$, No. 1, $P\bar{1}$, No. 2, and $P\bar{6}$, No. 174, respectively. The sequence of the graphs is

⁵ The HM symbols used here are nonconventional. They display the setting of the point group and follow the rules of *IT A*, Section 2.2.4.

2. MAXIMAL SUBGROUPS OF THE PLANE GROUPS AND SPACE GROUPS

determined by the sequence of the point groups in *IT A*, Table 2.1.2.1, fourth column. The graphs of $\bar{4}$, $\bar{3}$ and $6/m$ are nearly trivial, because to these crystal classes only two space-group types belong. The graphs of $mm2$ with 22, of mmm with 28 and of $4/mmm$ with 20 space-group types are the most complicated ones.

Isomorphic subgroups are special cases of k -subgroups. With the exception of both partners of the enantiomorphic space-group types, isomorphic subgroups are not displayed in the graphs. The explicit display of the isomorphic subgroups would add an infinite number of lines from each field for a space group back to this field, or at least one line (e.g. a circle) implicitly representing the infinite number of isomorphic subgroups, see the tables of maximal subgroups of Chapter 2.3.⁶ Such a line would have to be attached to every space-group symbol. Thus, there would be no more information.

Nevertheless, connections between isomorphic space groups are included indirectly if the group–subgroup chain encloses a space group of another type. In this case, a space group \mathcal{X} may be a subgroup of a space group \mathcal{Y} and \mathcal{Y}' a subgroup of \mathcal{X} , where \mathcal{Y} and \mathcal{Y}' belong to the same space-group type. The subgroup chain is then $\mathcal{Y} - \mathcal{X} - \mathcal{Y}'$. The two space groups \mathcal{Y} and \mathcal{Y}' are not identical but isomorphic. Whereas in general the label for the subgroup is positioned at a lower level than that for the original space group, for such relations the symbols for \mathcal{X} and \mathcal{Y} can only be drawn on the same level, connected by horizontal lines. If this happens at the top of a graph, the top level is occupied by more than one symbol (the number of symbols in the top level is the same as the number of symmorphic space-group types of the crystal class).

Horizontal lines are drawn as left or right arrows depending on the kind of relation. The arrow is always directed from the supergroup to the subgroup. If the relation is two-sided, as is always the case for enantiomorphic space-group types, then the relation is displayed by a pair of horizontal lines, one of them formed by a right and the other by a left arrow. In the graph in Fig. 2.5.1.5 for crystal class $mm2$, the connections of $Pmm2$ with $Cmm2$ and with $Amm2$ are displayed by double-headed arrows instead. Furthermore, some arrows in Fig. 2.5.1.5, crystal class $mm2$, and Fig. 2.5.1.6, mmm , are dashed or dotted in order to better distinguish the different lines and to increase clarity.

The different kinds of relations are demonstrated in the following examples.

Example 2.1.7.3.1.

In the graph in Fig. 2.5.1.1, crystal class 2, a space group $P2$ may be a subgroup of index 2 of a space group $C2$ by ‘Loss of centring translations’. On the other hand, subgroups of $P2$ in the block ‘Enlarged unit cell’, $\mathbf{a}' = 2\mathbf{a}$, $\mathbf{b}' = 2\mathbf{b}$, $C2(3)$ belong to the type $C2$, see the tables of maximal subgroups in Chapter 2.3. Therefore, both symbols are drawn at the same level and are connected by a pair of arrows pointing in opposite directions. Thus, the top level is occupied twice. In the graph in Fig. 2.5.1.2 of crystal class m , both the top level and the bottom level are occupied twice.

Example 2.1.7.3.2.

There are four symbols at the top level of the graph in Fig. 2.5.1.4, crystal class 222. Their relations are rather complicated. Whereas one can go (by index 2) from $P222$ directly to a subgroup of type $C222$ and *vice versa*, the connection from $F222$

directly to $C222$ is one-way. One always has to pass $C222$ on the way from $F222$ to a subgroup of the types $P222$ or $I222$. Thus, the only maximal subgroup of $F222$ among these groups is $C222$. One can go directly from $P222$ to $F222$ but not to $I222$ etc.

Because of the horizontal connecting arrows, it is clear that there cannot be much correspondence between the level in the graphs and the subgroup index. However, in no graph is a subgroup positioned at a higher level than the supergroup.

Example 2.1.7.3.3.

Consider the graph in Fig. 2.5.1.6 for crystal class mmm . To the space group $Cmmm$ (65) belong maximal non-isomorphic subgroups of the 11 space-group types (from left to right) $Ibam$ (72), $Cmcm$ (63), $Imma$ (74), $Pmnm$ (59), $Pbam$ (55), $Pban$ (50), $Pmma$ (51), $Pmna$ (53), $Cccm$ (66), $Pmmm$ (47) and $Immm$ (71). Although all of them have index 2, their symbols are positioned at very different levels of the graph.

The table for the subgroups of $Cmmm$ in Chapter 2.3 lists 22 non-isomorphic k -subgroups of index 2, because some of the space-group types mentioned above are represented by two or four different subgroups. This multiplicity cannot be displayed by multiple lines because the density of the lines in some of the k -subgroup graphs does not permit this kind of presentation, e.g. for mmm . The multiplicity may be taken from the subgroup tables in Chapter 2.3, where each non-isomorphic subgroup is listed individually.

Consider the connections from $Cmmm$ (65) to $Pbam$ (55). There are among others: the direct connection of index 2, the connection of index 4 over $Ibam$ (72), the connection of index 8 over $Imma$ (74) and $Pmma$ (51). Thus, starting from the same space group of type $Cmmm$ one arrives at different space groups of the type $Pbam$ with different unit cells but all belonging to the same space-group type and thus represented by the same field of the graph.

The index of a k -subgroup is restricted by Lemma 1.2.8.2.3 and by additional conditions. For the following statements one has to note that enantiomorphic space groups are isomorphic.

- (1) A non-isomorphic maximal k -subgroup of an oblique, rectangular or tetragonal plane group or of a triclinic, monoclinic, orthorhombic or tetragonal space group always has index 2.
- (2) In general, a non-isomorphic maximal k -subgroup \mathcal{H} of a trigonal space group \mathcal{G} has index 3. Exceptions are the pairs $P3m1-P3c1$, $P31m-P31c$, $R3m-R3c$, $\bar{P}31m-\bar{P}31c$, $\bar{P}3m1-\bar{P}3c1$ and $\bar{R}3m-\bar{R}3c$ with space-group Nos. between 156 and 167. They have index 2.
- (3) A non-isomorphic maximal k -subgroup \mathcal{H} of a hexagonal space group has index 2 or 3.
- (4) A non-isomorphic maximal k -subgroup \mathcal{H} of a cubic space group \mathcal{G} has either index 2 or index 4. The index is 2 if \mathcal{G} has an I lattice and \mathcal{H} a P lattice or if \mathcal{G} has a P lattice and \mathcal{H} an F lattice. The index is 4 if \mathcal{G} has an F lattice and \mathcal{H} a P lattice or if \mathcal{G} has a P lattice and \mathcal{H} an I lattice.

2.1.7.4. Graphs for plane groups

There are no graphs for plane groups in this volume. The four graphs for t -subgroups of plane groups are apart from the symbols the same as those for the corresponding space groups: $p4mm-P4mm$, $p6mm-P6mm$, $p2mg-Pma2$ and $p2gg-Pba2$, where the graphs for the space groups are included in the t -subgroup graphs in Figs. 2.4.1.1, 2.4.3.1, 2.4.2.1 and 2.4.2.3, respectively.

⁶ One could contemplate adding one line for each series of maximal isomorphic subgroups. However, the number of series depends on the rules that define the distribution of the isomorphic subgroups into the series and is thus not constant.

2.1. GUIDE TO THE SUBGROUP TABLES AND GRAPHS

The k -subgroup graphs are trivial for the plane groups $p1$, $p2$, $p4$, $p3$, $p6$ and $p6mm$ because there is only one plane group in its crystal class. The graphs for the crystal classes $4mm$ and $3m$ consist of two plane groups each: $p4mm$ and $p4gm$, $p3m1$ and $p31m$. Nevertheless, the graphs are different: the relation is one-sided for the tetragonal plane-group pair as it is in the space-group pair $P6/m$ (175)– $P6_3/m$ (176) and it is two-sided for the hexagonal plane-group pair as it is in the space-group pair $P\bar{4}$ (81)– $\bar{I}\bar{4}$ (82). The graph for the three plane groups of the crystal class m corresponds to the space-group graph for the crystal class 2.

Finally, the graph for the four plane groups of crystal class $2mm$ has no direct analogue among the k -subgroup graphs of the space groups. It can be obtained, however, from the graph in Fig. 2.5.1.3 of crystal class $2/m$ by removing the fields of $C2/c$ (15) and $P2_1/m$ (11) with all their connections to the remaining fields. The replacements are then: $C2/m$ (12) by $c2mm$ (9), $P2/m$ (10) by $p2mm$ (6), $P2/c$ (13) by $p2mg$ (7) and $P2_1/c$ (14) by $p2gg$ (8).

2.1.7.5. Application of the graphs

If a subgroup is not maximal then there must be a group–subgroup chain $\mathcal{G}-\mathcal{H}$ of maximal subgroups with more than two members which connects \mathcal{G} with \mathcal{H} . There are three possibilities: \mathcal{H} may be a t -subgroup or a k -subgroup or a general subgroup of \mathcal{G} . In the first two cases, the application of the graphs is straightforward because at least one of the graphs will permit one to find the possible chains directly. If \mathcal{H} is a k -subgroup of \mathcal{G} , isomorphic subgroups have to be included if necessary. If \mathcal{H} is a general subgroup of \mathcal{G} one has to combine t - and k -subgroup graphs, but the problem is only slightly more complicated. This is because for a general subgroup $\mathcal{H} < \mathcal{G}$, Hermann's theorem 1.2.8.1.2 states the existence of an intermediate group \mathcal{M} with $\mathcal{H} < \mathcal{M} < \mathcal{G}$ and the properties $\mathcal{H} < \mathcal{M}$ is a k -subgroup of \mathcal{M} and $\mathcal{M} < \mathcal{G}$ is a t -subgroup of \mathcal{G} .

Thus, however long and complicated the real chain may be, there is also always a chain for which only two graphs are needed: a t -subgroup graph for the relation between \mathcal{G} and \mathcal{M} and a k -subgroup graph for the relation between \mathcal{M} and \mathcal{H} .




Fig. 2.1.7.1. Contracted graph of the group–subgroup chains from $Fm\bar{3}m$ (225) to those subgroups with index 12 which belong to the space-group type $C2/m$ (12). The graph forms part of the total contracted graph of t -subgroups of $Fm\bar{3}m$ (Fig. 2.4.1.5).




Fig. 2.1.7.2. Complete graph of the group–subgroup chains from $Fm\bar{3}m$ (225) to one representative of those six $C2/m$ (12) subgroups with index 12 whose monoclinic axes are along the $\langle 110 \rangle$ directions of $Fm\bar{3}m$.

There is, however, a severe shortcoming to using contracted graphs for the analysis of group–subgroup relations, and great care has to be taken in such investigations. All subgroups \mathcal{H}_j with the same space-group type are represented by the same field of the graph, but these different non-maximal subgroups may permit different routes to a common original (super)group.

Example 2.1.7.5.1.

An example for *translationengleiche* subgroups is provided by the group–subgroup chain $Fm\bar{3}m$ (225)– $C2/m$ (12) of index 12. The contracted graph may be drawn by the program *Subgroupgraph* from the Bilbao Crystallographic Server, <http://www.cryst.ehu.es/>. It is shown in Fig. 2.1.7.1; each field represents all occurring subgroups of a space-group type: $I4/mmm$ (139) represents three subgroups, $R\bar{3}m$ (166) represents four subgroups, ... and $C2/m$ (12) represents nine subgroups belonging to two conjugacy classes. Fig. 2.1.7.1 is part of the contracted total graph of the *translationengleiche* subgroups of the space group $Fm\bar{3}m$, which is displayed in Fig.




Fig. 2.1.7.3. Complete graph of the group–subgroup chains from $Fm\bar{3}m$ (225) to one representative of those three $C2/m$ (12) subgroups with index 12 whose monoclinic axes are along the $\langle 001 \rangle$ directions of $Fm\bar{3}m$.

2. MAXIMAL SUBGROUPS OF THE PLANE GROUPS AND SPACE GROUPS

2.4.1.5. With *Subgroupgraph* one can also obtain the *complete graph* between $Fm\bar{3}m$ and the set of all nine subgroups of the type $C2/m$. It is too large to be reproduced here.

More instructive are the complete graphs for different single subgroups of the type $C2/m$ of $Fm\bar{3}m$. They can be obtained with the program *Symmodes* from the same server as above. In Fig. 2.1.7.2 such a graph is displayed for one of the six subgroups of type $C2/m$ of index 12 whose monoclinic axes point in the $\langle 110 \rangle$ directions of $Fm\bar{3}m$. Similarly, in Fig. 2.1.7.3 the complete graph is drawn for one of the three subgroups of $C2/m$ of index 12 whose monoclinic axes point in the $\langle 001 \rangle$ directions of $Fm\bar{3}m$. It differs markedly from the contracted graph and from the first complete graph. It is easily seen that it may be very misleading to use the contracted graph or the wrong individual complete graph instead of the right individual complete graph.

The following example deals with general subgroups.

Example 2.1.7.5.2.

The crystal structures of $SrTiO_3$ and $KCuF_3$ both belong to the space-group type $I4/mcm$, No. 140. They can be derived by different distortions from the same ideal perovskite ABX_3 structure with the space group $Pm\bar{3}m$, No. 221. Common to both chains is Hermann's group \mathcal{M} of space group $P4/mmm$, No. 123, and the unit cell of ABX_3 . This is the only intermediate group for $SrTiO_3$. For the pair ABX_3 – $KCuF_3$, on the other hand, there exists another chain with an intermediate space group of type $Fm\bar{3}c$, No. 226. The combination of the graph in Fig. 2.4.1.1 for t -subgroups with the graph in Fig. 2.5.2.7 for k -subgroups is thus possible for both crystal–chemical relations. The combination of the graph in Fig. 2.5.5.5 for k -subgroups with the graph in Fig. 2.4.1.6 for t -subgroups is, however, only meaningful for the chain ABX_3 – $KCuF_3$. This cannot be concluded from the contracted graphs but can be seen from the complete graph, as displayed in Fig. 1 of Wondratschek & Aroyo (2001).