

C_{2v}^4
 $Pma2$

No. 28

 $Pma2$
Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3)

General position

 Multiplicity,
 Wyckoff letter,
 Site symmetry

Coordinates

 4 d 1 (1) x, y, z (2) \bar{x}, \bar{y}, z (3) $x + \frac{1}{2}, \bar{y}, z$ (4) $\bar{x} + \frac{1}{2}, y, z$
I Maximal translationengleiche subgroups

[2] $P1a1$ (7, $P1c1$)	1; 3	$-\mathbf{a} - \mathbf{c}, \mathbf{b}, \mathbf{a}$	
[2] $Pm11$ (6, $P1m1$)	1; 4	$\mathbf{c}, \mathbf{a}, \mathbf{b}$	1/4, 0, 0
[2] $P112$ (3)	1; 2		

II Maximal klassengleiche subgroups

• Enlarged unit cell

[2] $\mathbf{b}' = 2\mathbf{b}$			
$Pba2$ (32)	$\langle 2; 3 + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	
$Pba2$ (32)	$\langle 3; 2 + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	0, 1/2, 0
$Pma2$ (28)	$\langle 2; 3 \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	
$Pma2$ (28)	$\langle (2; 3) + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, \mathbf{c}$	0, 1/2, 0
[2] $\mathbf{c}' = 2\mathbf{c}$			
$Pmn2_1$ (31)	$\langle (2; 3) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	1/4, 0, 0
$Pcn2$ (30, $Pnc2$)	$\langle 2; 3 + (0, 0, 1) \rangle$	$-\mathbf{b}, \mathbf{a}, 2\mathbf{c}$	
$Pca2_1$ (29)	$\langle 3; 2 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	
$Pma2$ (28)	$\langle 2; 3 \rangle$	$\mathbf{a}, \mathbf{b}, 2\mathbf{c}$	
[2] $\mathbf{b}' = 2\mathbf{b}, \mathbf{c}' = 2\mathbf{c}$			
$Aea2$ (41)	$\langle 2; 3 + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	
$Aea2$ (41)	$\langle 3; 2 + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	0, 1/2, 0
$Ama2$ (40)	$\langle 2; 3 \rangle$	$\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	
$Ama2$ (40)	$\langle (2; 3) + (0, 1, 0) \rangle$	$\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	0, 1/2, 0
[3] $\mathbf{a}' = 3\mathbf{a}$			
$Pma2$ (28)	$\langle 2; 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	
$Pma2$ (28)	$\langle 2 + (2, 0, 0); 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	1, 0, 0
$Pma2$ (28)	$\langle 2 + (4, 0, 0); 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{b}' = 3\mathbf{b}$			
$Pma2$ (28)	$\langle 2; 3 \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	
$Pma2$ (28)	$\langle (2; 3) + (0, 2, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 1, 0
$Pma2$ (28)	$\langle (2; 3) + (0, 4, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 2, 0
[3] $\mathbf{c}' = 3\mathbf{c}$			
$Pma2$ (28)	$\langle 2; 3 \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	

• Series of maximal isomorphic subgroups

[p] $\mathbf{a}' = p\mathbf{a}$			
$Pma2$ (28)	$\langle 2 + (2u, 0, 0); 3 + (\frac{p}{2} - \frac{1}{2}, 0, 0) \rangle$	$p\mathbf{a}, \mathbf{b}, \mathbf{c}$	$u, 0, 0$
	$p > 2; 0 \leq u < p$		
	p conjugate subgroups for the prime p		
[p] $\mathbf{b}' = p\mathbf{b}$			
$Pma2$ (28)	$\langle (2; 3) + (0, 2u, 0) \rangle$	$\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$0, u, 0$
	$p > 2; 0 \leq u < p$		
	p conjugate subgroups for the prime p		
[p] $\mathbf{c}' = p\mathbf{c}$			
$Pma2$ (28)	$\langle 2; 3 \rangle$	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	
	$p > 1$		
	no conjugate subgroups		

I Minimal translationengleiche supergroups

 [2] $Pccm$ (49); [2] $Pmma$ (51); [2] $Pmna$ (53); [2] $Pbcm$ (57)

II Minimal non-isomorphic klassengleiche supergroups

• Additional centring translations

 [2] $Cmm2$ (35); [2] $Bme2$ (39, $Aem2$); [2] $Ama2$ (40); [2] $Ima2$ (46)

• Decreased unit cell

 [2] $\mathbf{a}' = \frac{1}{2}\mathbf{a}$ $Pmm2$ (25)