

C_{4v}^3
 $P4_2cm$

No. 101

 $P4_2cm$
Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3); (5)

General position

 Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

8	<i>e</i>	1		(1) x, y, z	(2) \bar{x}, \bar{y}, z	(3) $\bar{y}, x, z + \frac{1}{2}$	(4) $y, \bar{x}, z + \frac{1}{2}$
				(5) $x, \bar{y}, z + \frac{1}{2}$	(6) $\bar{x}, y, z + \frac{1}{2}$	(7) \bar{y}, \bar{x}, z	(8) y, x, z

I Maximal translationengleiche subgroups

[2] $P4_211$ (77, $P4_2$)	1; 2; 3; 4	
[2] $P21m$ (35, $Cmm2$)	1; 2; 7; 8	a – b, a + b, c
[2] $P2c1$ (27, $Pcc2$)	1; 2; 5; 6	

II Maximal klassengleiche subgroups

• Enlarged unit cell

[2] $\mathbf{a}' = 2\mathbf{a}, \mathbf{b}' = 2\mathbf{b}$		
$C4_2cd$ (106, $P4_2bc$)	$\langle 2; 3; 5 + (0, 1, 0) \rangle$	a – b, a + b, c
$C4_2cd$ (106, $P4_2bc$)	$\langle 2; 5; 3 + (1, 0, 0) \rangle$	a – b, a + b, c 1/2, 1/2, 0
$C4_2cm$ (105, $P4_2mc$)	$\langle 2; 3; 5 \rangle$	a – b, a + b, c
$C4_2cm$ (105, $P4_2mc$)	$\langle 2; 3 + (1, 0, 0); 5 + (0, 1, 0) \rangle$	a – b, a + b, c 1/2, 1/2, 0
[3] $\mathbf{c}' = 3\mathbf{c}$		
$P4_2cm$ (101)	$\langle 2; (3; 5) + (0, 0, 1) \rangle$	a, b, 3c

• Series of maximal isomorphic subgroups

[<i>p</i>] $\mathbf{c}' = p\mathbf{c}$		
$P4_2cm$ (101)	$\langle 2; (3; 5) + (0, 0, \frac{p}{2} - \frac{1}{2}) \rangle$ $p > 2$ no conjugate subgroups	a, b, pc
[p^2] $\mathbf{a}' = p\mathbf{a}, \mathbf{b}' = p\mathbf{b}$		
$P4_2cm$ (101)	$\langle 2 + (2u, 2v, 0); 3 + (u + v, -u + v, 0); 5 + (0, 2v, 0) \rangle$ $p > 2; 0 \leq u < p; 0 \leq v < p$ p^2 conjugate subgroups for the prime p	pa, pb, c $u, v, 0$

I Minimal translationengleiche supergroups

 [2] $P4_2/mcm$ (132); [2] $P4_2/ncm$ (138)

II Minimal non-isomorphic klassengleiche supergroups

• Additional centring translations

 [2] $C4_2cm$ (105, $P4_2mc$); [2] $I4cm$ (108)

• Decreased unit cell

 [2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $P4mm$ (99)