

1.2. GENERAL INTRODUCTION TO THE SUBGROUPS OF SPACE GROUPS

Example 1.2.6.2.4

Consider a space group \mathcal{G} of the type $C121$, referred to a conventional coordinate system. The set $\mathcal{T}(\mathcal{G})$ of all translations can be split into the set \mathcal{T}_i of all translations with integer coefficients u, v and w and the set \mathcal{T}_f of all translations for which the coefficients u and v are fractional. The set \mathcal{T}_i forms a group; the set \mathcal{T}_f is the other coset in the decomposition $(\mathcal{T}(\mathcal{G}) : \mathcal{T}_i)$ and does not form a group. Let t_C be the ‘centring translation’ with the translation vector $\frac{1}{2}(\mathbf{a} + \mathbf{b})$. Then \mathcal{T}_f can be written $\mathcal{T}_i t_C$. Let $\mathbf{2}_0$ mean a twofold rotation around the rotation axis through the origin. There are altogether four cosets of the decomposition $(\mathcal{G} : \mathcal{T}_i)$, which can be written now as $\mathcal{T}_i, \mathcal{T}_f = \mathcal{T}_i t_C, \mathcal{T}_i \mathbf{2}_0$ and $\mathcal{T}_f \mathbf{2}_0 = (\mathcal{T}_i t_C) \mathbf{2}_0 = \mathcal{T}_i (t_C \mathbf{2}_0)$. The union $\mathcal{T}_i \cup (\mathcal{T}_i t_C) = \mathcal{T}_{\mathcal{G}}$ forms the *translationengleiche* maximal subgroup $C1$ (conventional setting $P1$) of \mathcal{G} of index 2. The union $\mathcal{T}_i \cup (\mathcal{T}_i \mathbf{2}_0)$ forms the *klassengleiche* maximal subgroup $P121$ of \mathcal{G} of index 2. The union $\mathcal{T}_i \cup (\mathcal{T}_i (t_C \mathbf{2}_0))$ also forms a *klassengleiche* maximal subgroup of index 2. Its HM symbol is $P12_1$, and the twofold rotations $\mathbf{2}$ of the point group $\mathbf{2}$ are realized by screw rotations $\mathbf{2}_1$ in this subgroup because $(t_C \mathbf{2}_0)$ is a screw rotation with its screw axis running parallel to the \mathbf{b} axis through the point $\frac{1}{4}, 0, 0$. There are in fact these two k -subgroups of $C121$ of index 2 which have the group \mathcal{T}_i in common. In the subgroup table of $C121$ both are listed under the heading ‘Loss of centring translations’ because the conventional unit cell is retained while only the centring translations have disappeared. (Four additional *klassengleiche* maximal subgroups of $C121$ with index 2 are found under the heading ‘Enlarged unit cell’.)

The group \mathcal{T}_i of type $P1$ is a non-maximal subgroup of $C121$ of index 4.

Definition 1.2.6.2.5. A *klassengleiche* or k -subgroup $\mathcal{H} < \mathcal{G}$ is called *isomorphic* or an *isomorphic subgroup* if it belongs to the same affine space-group type (isomorphism type) as \mathcal{G} . If a subgroup is not isomorphic, it is sometimes called *non-isomorphic*. \square

Isomorphic subgroups are special k -subgroups. The importance of the distinction between k -subgroups in general and isomorphic subgroups in particular stems from the fact that the number of maximal non-isomorphic k -subgroups of any space group is finite, whereas the number of maximal isomorphic subgroups is always infinite, see Section 1.2.8.

Example 1.2.6.2.6

Consider a space group \mathcal{G} of type $P\bar{1}$ referred to a conventional coordinate system. The translation subgroup $\mathcal{T}(\mathcal{G})$ consists of all translations with translation vectors $\mathbf{t} = u\mathbf{a} + v\mathbf{b} + w\mathbf{c}$, where u, v and w run through all integer numbers. There is an inversion $\bar{1}_0$ with the inversion point at the origin and also an infinite number of other inversions, generated by the combinations of $\bar{1}_0$ with all translations of $\mathcal{T}(\mathcal{G})$.

We consider the subgroup $\mathcal{T}_{\mathcal{G}}$ of all translations with an even coefficient u and arbitrary integers v and w as well as the coset decomposition $(\mathcal{G} : \mathcal{T}_{\mathcal{G}})$. Let t_a be the translation with the translation vector \mathbf{a} . There are four cosets: $\mathcal{T}_{\mathcal{G}}, \mathcal{T}_{\mathcal{G}} t_a, \mathcal{T}_{\mathcal{G}} \bar{1}_0$ and $\mathcal{T}_{\mathcal{G}} (t_a \bar{1}_0)$. The union $\mathcal{T}_{\mathcal{G}} \cup (\mathcal{T}_{\mathcal{G}} t_a)$ forms the *translationengleiche* maximal subgroup $\mathcal{T}(\mathcal{G})$ of index 2. The union $\mathcal{T}_{\mathcal{G}} \cup (\mathcal{T}_{\mathcal{G}} \bar{1}_0)$ forms an isomorphic maximal subgroup of index 2, as does the union $\mathcal{T}_{\mathcal{G}} \cup (\mathcal{T}_{\mathcal{G}} (t_a \bar{1}_0))$. There are thus two maximal isomorphic subgroups of index 2 which are obtained by doubling the a lattice parameter. There are altogether 14 isomorphic sub-

groups of index 2 for any space group of type $P\bar{1}$ which are obtained by seven different cell enlargements.

If \mathcal{G} belongs to a pair of enantiomorphic space-group types, then the isomorphic subgroups of \mathcal{G} may belong to different crystallographic space-group types with different HM symbols and different space-group numbers. In this case, an infinite number of subgroups belong to the crystallographic space-group type of \mathcal{G} and another infinite number belong to the enantiomorphic space-group type.

Example 1.2.6.2.7

Space group $P4_1$, No. 76, has for any prime number $p > 2$ an isomorphic maximal subgroup of index p with the lattice parameters a, b, pc . This is an infinite number of subgroups because there is an infinite number of primes. The subgroups belong to the space-group type $P4_1$ if $p = 4n + 1$; they belong to the type $P4_3$ if $p = 4n + 3$.

Definition 1.2.6.2.8. A subgroup of a space group is called *general* or a *general subgroup* if it is neither a *translationengleiche* nor a *klassengleiche* subgroup. It has lost translations as well as linear parts, i.e. point-group symmetry. \square

Example 1.2.6.2.9

The subgroup $\mathcal{T}_{\mathcal{G}}$ in Example 1.2.6.2.6 has lost all inversions of the original space group $P\bar{1}$ as well as all translations with odd u . It is a general subgroup $P1$ of the space group $P\bar{1}$ of index 4.

1.2.6.3. The role of normalizers for group–subgroup pairs of space groups

In Section 1.2.4.5, the normalizer $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$ of a subgroup $\mathcal{H} < \mathcal{G}$ in the group \mathcal{G} was defined. The equation $\mathcal{H} \trianglelefteq \mathcal{N}_{\mathcal{G}}(\mathcal{H}) \leq \mathcal{G}$ holds, i.e. \mathcal{H} is a normal subgroup of $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$. The normalizer $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$, by its index in \mathcal{G} , determines the number $N_j = |\mathcal{G} : \mathcal{N}_{\mathcal{G}}(\mathcal{H})|$ of subgroups $\mathcal{H}_j < \mathcal{G}$ that are conjugate in the group \mathcal{G} , cf. Remarks (2) and (3) below Definition 1.2.4.5.1.

The group–subgroup relations between space groups become more transparent if one looks at them from a more general point of view. Space groups are part of the general theory of mappings. Particular groups are the *affine group* \mathcal{A} of all reversible affine mappings, the *Euclidean group* \mathcal{E} of all isometries, the *translation group* \mathcal{T} of all translations and the *orthogonal group* \mathcal{O} of all orthogonal mappings.

Connected with any particular space group \mathcal{G} are its group of translations $\mathcal{T}(\mathcal{G})$ and its point group $\mathcal{P}_{\mathcal{G}}$. In addition, the normalizers $\mathcal{N}_{\mathcal{A}}(\mathcal{G})$ of \mathcal{G} in the affine group \mathcal{A} and $\mathcal{N}_{\mathcal{E}}(\mathcal{G})$ in the Euclidean group \mathcal{E} are useful. They are listed in Section 15.2.1 of IT A. Although consisting of isometries only, $\mathcal{N}_{\mathcal{E}}(\mathcal{G})$ is not necessarily a space group, see the second paragraph of Example 1.2.7.3.1.

For the group–subgroup pairs $\mathcal{H} < \mathcal{G}$ the following relations hold:

- (1) $\mathcal{T}(\mathcal{H}) \leq \mathcal{H} \leq \mathcal{N}_{\mathcal{G}}(\mathcal{H}) \leq \mathcal{G} \leq \mathcal{N}_{\mathcal{E}}(\mathcal{G}) < \mathcal{E}$;
 (1a) $\mathcal{H} \leq \mathcal{N}_{\mathcal{G}}(\mathcal{H}) \leq \mathcal{N}_{\mathcal{E}}(\mathcal{H}) < \mathcal{E}$;
 (1b) $\mathcal{N}_{\mathcal{E}}(\mathcal{H}) \leq \mathcal{N}_{\mathcal{A}}(\mathcal{H}) < \mathcal{A}$;
- (2) $\mathcal{T}(\mathcal{H}) \leq \mathcal{T}(\mathcal{G}) < \mathcal{T} < \mathcal{E}$;
- (3) $\mathcal{T}(\mathcal{G}) \leq \mathcal{G} \leq \mathcal{N}_{\mathcal{E}}(\mathcal{G}) \leq \mathcal{N}_{\mathcal{A}}(\mathcal{G}) < \mathcal{A}$.

The subgroup \mathcal{H} may be a *translationengleiche* or a *klassengleiche* or a general subgroup of \mathcal{G} . In any case, the normalizer

1. SPACE GROUPS AND THEIR SUBGROUPS

$\mathcal{N}_{\mathcal{G}}(\mathcal{H})$ determines the length of the conjugacy class of $\mathcal{H} < \mathcal{G}$, but it is not feasible to list for each group–subgroup pair $\mathcal{H} < \mathcal{G}$ its normalizer $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$. Indeed, it is only necessary to list for any space group \mathcal{H} its normalizer $\mathcal{N}_{\mathcal{E}}(\mathcal{H})$ in the Euclidean group \mathcal{E} of all isometries, as is done in *IT A*, Section 15.2.1. From such a list the normalizers for the group–subgroup pairs can be obtained easily, because for any chain of space groups $\mathcal{H} < \mathcal{G} < \mathcal{E}$, the relations $\mathcal{H} \leq \mathcal{N}_{\mathcal{G}}(\mathcal{H}) \leq \mathcal{G}$ and $\mathcal{H} \leq \mathcal{N}_{\mathcal{G}}(\mathcal{H}) \leq \mathcal{N}_{\mathcal{E}}(\mathcal{H})$ hold. The normalizer $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$ consists consequently of all those isometries of $\mathcal{N}_{\mathcal{E}}(\mathcal{H})$ that are also elements of \mathcal{G} , *i.e.* that belong to the intersection $\mathcal{N}_{\mathcal{E}}(\mathcal{H}) \cap \mathcal{G}$, *cf.* the examples of Section 1.2.7.¹¹

The isomorphism type of the Euclidean normalizer $\mathcal{N}_{\mathcal{E}}(\mathcal{H})$ may depend on the lattice parameters of the space group (*specialized* Euclidean normalizer). For example, if the lattice of the space group $P\bar{1}$ of a triclinic crystal is accidentally monoclinic at a certain temperature and pressure or for a certain composition in a continuous solid-solution series, then the Euclidean normalizer of this space group belongs to the space-group types $P2/m$ or $C2/m$, otherwise it belongs to $P\bar{1}$. Such a *specialized Euclidean normalizer* (here $P2/m$ or $C2/m$) may be distinguished from the *typical Euclidean normalizer* (here $P\bar{1}$), for which the lattice of \mathcal{H} is not more symmetric than is required by the symmetry of \mathcal{H} . The specialized Euclidean normalizers were first listed in the 5th edition of *IT A* (2005), Section 15.2.1.

1.2.7. Application to domain structures

1.2.7.1. Introductory remarks

In this section, the basic group-theoretical aspects of this chapter are exemplified using the topic of domain structures (transformation twins). Domain structures result from a displacive or order–disorder phase transition. A homogeneous single crystal phase **A** (parent or prototypic phase) is transformed to a crystalline phase **B** (daughter phase, distorted phase). In most cases phase **B** is inhomogeneous, consisting of homogeneous regions which are called *domains*.

Definition 1.2.7.1.1. A connected homogeneous part of a domain structure or of a twinned crystal of phase **B** is called a *domain*. Each domain is a single crystal. The part of space that is occupied by a domain is the *region* of that domain. \square

The space groups \mathcal{H}_j of phase **B** are conjugate subgroups of the space group \mathcal{G} of phase **A**, $\mathcal{H}_j < \mathcal{G}$. The number of domains is not limited; they differ in their locations in space, in their orientations, in their sizes, in their shapes and in their space groups $\mathcal{H}_j < \mathcal{G}$ which, however, all belong to the same space-group type \mathcal{H} . The boundaries between the domains, called the domain walls, are assumed to be (infinitely) thin.

A deeper discussion of domain structures or transformation twins and their properties needs a much more detailed treatment, as is given in Volume D of *International Tables for Crystallography* (2003) (abbreviated as *IT D*) Part 3, by Janovec, Hahn & Klapper (Chapter 3.2), by Hahn & Klapper (Chapter 3.3) and by Janovec & Přívratská (Chapter 3.4) with more than 400

references. Domains are also considered in Section 1.6.6 of this volume.

In this section, non-ferroelastic phase transitions are treated without any special assumption as well as ferroelastic phase transitions under the simplifying *parent clamping approximation*, abbreviated PCA, introduced by Janovec *et al.* (1989), see also *IT D*, Section 3.4.2.5. A transition is *non-ferroelastic* if the strain tensors (metric tensors) of the low-symmetry phase **B** have the same independent components as the strain tensor of the phase **A**.¹² There are thus *no spontaneous strain components* which distort the lattices of the domains. In a *ferroelastic* phase transition the strain tensors of phase **B** have more independent components than the strain tensor of phase **A**. The additional strain components cause lattice strain. By the PCA the lattice parameters of phase **B** at the transition are adapted to those of phase **A**, *i.e.* to the lattice symmetry of phase **A**. Therefore, under the PCA the ferroelastic phases display the same behaviour as the non-ferroelastic phases.

If in this section ferroelastic phase transitions are considered, the PCA is assumed to be applied.

Under a non-ferroelastic phase transition or under the assumption of the PCA, the translations of the constituents of the phase **B** are translations of phase **A** and the space groups \mathcal{H}_j of **B** are subgroups of the space group \mathcal{G} of **A**, $\mathcal{H}_j < \mathcal{G}$.

Under this supposition the domain structure formed may exhibit different chiralities and/or polarities of its domains with different spatial orientations of their symmetry elements. Nevertheless, each domain has the same specific energy and the lattice of each domain is part of the lattice of the parent structure **A** with space group \mathcal{G} .

The description of domain structures by their crystal structures is called the *microscopic description*, *IT D*, Section 3.4.2.1. In the *continuum description*, the crystals are treated as anisotropic continua, *IT D*, Section 3.4.2.1. The role of the space groups is then taken over by the point groups of the domains. The continuum description is used when one is essentially interested in the macroscopic physical properties of the domain structure.

Different kinds of nomenclature are used in the discussion of domain structures. The basic concepts of *domain* and *domain state* are established in *IT D*, Section 3.4.2.1; the terms *symmetry state* and *directional state* are newly introduced here in the context of domain structures. All these concepts classify the domains and will be defined in the next section and applied in different examples of phase transitions in Section 1.2.7.3.

1.2.7.2. Domain states, symmetry states and directional states

In order to describe what happens in a phase transition of the kind considered, a few notions are useful.

If the domains of phase **B** have been formed from a single crystal of phase **A**, then they belong to a finite (small) number of domain states \mathbf{B}_j with space groups \mathcal{H}_j . The domain states have well defined relations to the original crystal of phase **A** and its space group \mathcal{G} . In order to describe these relations, the notion of crystal pattern is used. Any perfect (ideal) crystal is a finite block of the corresponding infinite arrangement, the symmetry of which is a space group and thus contains translations. Here, this

¹¹ For *maximal* subgroups, a calculation of the conjugacy classes is not necessary because these are indicated in the subgroup tables of Part 2 of this volume by braces to the left of the data sets for the low-index subgroups and by text for the series of isomorphic subgroups. For non-maximal subgroups, the conjugacy relations are not indicated but can be calculated in the way described here. They are also available online on the Bilbao Crystallographic Server, <http://www.cryst.ehu.es/>, under the program *Subgroupaph*.

¹² A phase transition is non-ferroelastic if the space groups \mathcal{G} of the high-symmetry phase **A** and \mathcal{H}_j of the low-symmetry phase **B** belong to the *same crystal family*, of which there are six: triclinic, monoclinic, orthorhombic, tetragonal, trigonal–hexagonal and cubic. In ferroelastic phase transitions the space groups \mathcal{G} of **A** and \mathcal{H}_j of **B** belong to different crystal families. Only then can spontaneous strain components occur. They are avoided by the assumption of the PCA.