

$C_{2v}^8$ 
 $Pba2$ 

No. 32

 $Pba2$ 
**Generators selected** (1);  $t(1,0,0)$ ;  $t(0,1,0)$ ;  $t(0,0,1)$ ; (2); (3)

**General position**

 Multiplicity,  
 Wyckoff letter,  
 Site symmetry

Coordinates

 4  $c$  1 (1)  $x, y, z$  (2)  $\bar{x}, \bar{y}, z$  (3)  $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z$  (4)  $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, z$ 
**I Maximal translationengleiche subgroups**

[2] $P1a1$ (7, $P1c1$ )	1; 3	$-a - c, b, a$	0, 1/4, 0
[2] $Pb11$ (7, $P1c1$ )	1; 4	$c, a, b$	1/4, 0, 0
[2] $P112$ (3)	1; 2		

**II Maximal klassengleiche subgroups**

## • Enlarged unit cell

[2] $c' = 2c$			
$Pnn2$ (34)	$\langle 2; 3 + (0, 0, 1) \rangle$	$a, b, 2c$	
$Pna2_1$ (33)	$\langle 3; 2 + (0, 0, 1) \rangle$	$a, b, 2c$	
$Pbn2_1$ (33, $Pna2_1$ )	$\langle (2; 3) + (0, 0, 1) \rangle$	$-b, a, 2c$	
$Pba2$ (32)	$\langle 2; 3 \rangle$	$a, b, 2c$	
[3] $a' = 3a$			
$Pba2$ (32)	$\langle 2; 3 + (1, 0, 0) \rangle$	$3a, b, c$	
$Pba2$ (32)	$\langle 2 + (2, 0, 0); 3 + (1, 0, 0) \rangle$	$3a, b, c$	1, 0, 0
$Pba2$ (32)	$\langle 2 + (4, 0, 0); 3 + (1, 0, 0) \rangle$	$3a, b, c$	2, 0, 0
[3] $b' = 3b$			
$Pba2$ (32)	$\langle 2; 3 + (0, 1, 0) \rangle$	$a, 3b, c$	
$Pba2$ (32)	$\langle 2 + (0, 2, 0); 3 + (0, 3, 0) \rangle$	$a, 3b, c$	0, 1, 0
$Pba2$ (32)	$\langle 2 + (0, 4, 0); 3 + (0, 5, 0) \rangle$	$a, 3b, c$	0, 2, 0
[3] $c' = 3c$			
$Pba2$ (32)	$\langle 2; 3 \rangle$	$a, b, 3c$	

## • Series of maximal isomorphic subgroups

[ $p$ ] $a' = pa$			
$Pba2$ (32)	$\langle 2 + (2u, 0, 0); 3 + (\frac{p}{2} - \frac{1}{2}, 0, 0) \rangle$ prime $p > 2$ ; $0 \leq u < p$ $p$ conjugate subgroups	$pa, b, c$	$u, 0, 0$
[ $p$ ] $b' = pb$			
$Pba2$ (32)	$\langle 2 + (0, 2u, 0); 3 + (0, \frac{p}{2} - \frac{1}{2} + 2u, 0) \rangle$ prime $p > 2$ ; $0 \leq u < p$ $p$ conjugate subgroups	$a, pb, c$	$0, u, 0$
[ $p$ ] $c' = pc$			
$Pba2$ (32)	$\langle 2; 3 \rangle$ $p$ prime no conjugate subgroups	$a, b, pc$	

**I Minimal translationengleiche supergroups**

 [2]  $Pban$  (50); [2]  $Pcca$  (54); [2]  $Pbam$  (55); [2]  $P4bm$  (100); [2]  $P4_2bc$  (106); [2]  $P\bar{4}b2$  (117)

**II Minimal non-isomorphic klassengleiche supergroups**

## • Additional centring translations

 [2]  $Cmm2$  (35); [2]  $Aea2$  (41); [2]  $Bbe2$  (41,  $Aea2$ ); [2]  $Iba2$  (45)

## • Decreased unit cell

 [2]  $a' = \frac{1}{2}a$   $Pbm2$  (28,  $Pma2$ ); [2]  $b' = \frac{1}{2}b$   $Pma2$  (28)