

C_{2v}^{10} $Pnn2$

No. 34

 $Pnn2$ Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3)

General position

Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

4 c 1 (1) x, y, z (2) \bar{x}, \bar{y}, z (3) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z + \frac{1}{2}$ (4) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, z + \frac{1}{2}$ I Maximal *translationengleiche* subgroups

[2] $P1n1$ (7, $P1c1$)	1; 3	$\mathbf{c}, \mathbf{b}, -\mathbf{a} - \mathbf{c}$	0, 1/4, 0
[2] $Pn11$ (7, $P1c1$)	1; 4	$\mathbf{b}, \mathbf{a}, -\mathbf{b} - \mathbf{c}$	1/4, 0, 0
[2] $P112$ (3)	1; 2		

II Maximal *klassengleiche* subgroups

• Enlarged unit cell

[2] $\mathbf{a}' = 2\mathbf{a}, \mathbf{b}' = 2\mathbf{b}, \mathbf{c}' = 2\mathbf{c}$

$Fdd2$ (43)	$\langle 2; 3 \rangle$	$2\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	
$Fdd2$ (43)	$\langle 3; 2 + (1, 0, 0) \rangle$	$2\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	1/2, 0, 0
$Fdd2$ (43)	$\langle (2; 3) + (0, 1, 0) \rangle$	$2\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	0, 1/2, 0
$Fdd2$ (43)	$\langle 2 + (1, 1, 0); 3 + (0, 1, 0) \rangle$	$2\mathbf{a}, 2\mathbf{b}, 2\mathbf{c}$	1/2, 1/2, 0
[3] $\mathbf{a}' = 3\mathbf{a}$			
$Pnn2$ (34)	$\langle 2; 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	
$Pnn2$ (34)	$\langle 2 + (2, 0, 0); 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	1, 0, 0
$Pnn2$ (34)	$\langle 2 + (4, 0, 0); 3 + (1, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{b}' = 3\mathbf{b}$			
$Pnn2$ (34)	$\langle 2; 3 + (0, 1, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	
$Pnn2$ (34)	$\langle 2 + (0, 2, 0); 3 + (0, 3, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 1, 0
$Pnn2$ (34)	$\langle 2 + (0, 4, 0); 3 + (0, 5, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 2, 0
[3] $\mathbf{c}' = 3\mathbf{c}$			
$Pnn2$ (34)	$\langle 2; 3 + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	

• Series of maximal isomorphic subgroups

[p] $\mathbf{a}' = p\mathbf{a}$

$Pnn2$ (34)	$\langle 2 + (2u, 0, 0); 3 + (\frac{p}{2} - \frac{1}{2}, 0, 0) \rangle$ prime $p > 2$; $0 \leq u < p$ p conjugate subgroups	$p\mathbf{a}, \mathbf{b}, \mathbf{c}$	$u, 0, 0$
-------------	--	---------------------------------------	-----------

[p] $\mathbf{b}' = p\mathbf{b}$

$Pnn2$ (34)	$\langle 2 + (0, 2u, 0); 3 + (0, \frac{p}{2} - \frac{1}{2} + 2u, 0) \rangle$ prime $p > 2$; $0 \leq u < p$ p conjugate subgroups	$\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$0, u, 0$
-------------	---	---------------------------------------	-----------

[p] $\mathbf{c}' = p\mathbf{c}$

$Pnn2$ (34)	$\langle 2; 3 + (0, 0, \frac{p}{2} - \frac{1}{2}) \rangle$ prime $p > 2$ no conjugate subgroups	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	
-------------	---	---------------------------------------	--

I Minimal *translationengleiche* supergroups[2] $Pnnn$ (48); [2] $Pnna$ (52); [2] $Pnmm$ (58); [2] $P4_2nm$ (102); [2] $P4nc$ (104); [2] $P\bar{4}n2$ (118)II Minimal non-isomorphic *klassengleiche* supergroups

• Additional centring translations

[2] $Ccc2$ (37); [2] $Ama2$ (40); [2] $Bbm2$ (40, $Ama2$); [2] $Imm2$ (44)

• Decreased unit cell

[2] $\mathbf{a}' = \frac{1}{2}\mathbf{a}$ $Pnc2$ (30); [2] $\mathbf{b}' = \frac{1}{2}\mathbf{b}$ $Pcn2$ (30, $Pnc2$); [2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $Pba2$ (32)