

C_{2v}^{12}
 $Cmc2_1$

No. 36

 $Cmc2_1$
Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{1}{2},\frac{1}{2},0)$; (2); (3)

General position

 Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

 $(0,0,0)+ (\frac{1}{2},\frac{1}{2},0)+$

 8 b 1

 (1) x,y,z (2) $\bar{x},\bar{y},z+\frac{1}{2}$ (3) $x,\bar{y},z+\frac{1}{2}$ (4) \bar{x},y,z
I Maximal translationengleiche subgroups

[2] $C1c1$ (9)	(1; 3)+	
[2] $Cm11$ (8, $C1m1$)	(1; 4)+	$-\mathbf{b}, \mathbf{a}, \mathbf{c}$
[2] $C112_1$ (4, $P112_1$)	(1; 2)+	$1/2(\mathbf{a}-\mathbf{b}), 1/2(\mathbf{a}+\mathbf{b}), \mathbf{c}$

II Maximal klassengleiche subgroups

• Loss of centring translations

[2] $Pbn2_1$ (33, $Pna2_1$)	1; 2; (3; 4) + $(\frac{1}{2}, \frac{1}{2}, 0)$	$-\mathbf{b}, \mathbf{a}, \mathbf{c}$	
[2] $Pmn2_1$ (31)	1; 4; (2; 3) + $(\frac{1}{2}, \frac{1}{2}, 0)$		0, 1/4, 0
[2] $Pbc2_1$ (29, $Pca2_1$)	1; 3; (2; 4) + $(\frac{1}{2}, \frac{1}{2}, 0)$	$-\mathbf{b}, \mathbf{a}, \mathbf{c}$	1/4, 1/4, 0
[2] $Pmc2_1$ (26)	1; 2; 3; 4		

• Enlarged unit cell

[3] $\mathbf{a}' = 3\mathbf{a}$			
$\left\{ \begin{array}{l} Cmc2_1 (36) \\ Cmc2_1 (36) \\ Cmc2_1 (36) \end{array} \right.$	$\langle 2; 3 \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	
	$\langle 3; 2 + (2, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	1, 0, 0
	$\langle 3; 2 + (4, 0, 0) \rangle$	$3\mathbf{a}, \mathbf{b}, \mathbf{c}$	2, 0, 0
[3] $\mathbf{b}' = 3\mathbf{b}$			
$\left\{ \begin{array}{l} Cmc2_1 (36) \\ Cmc2_1 (36) \\ Cmc2_1 (36) \end{array} \right.$	$\langle 2; 3 \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	
	$\langle (2; 3) + (0, 2, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 1, 0
	$\langle (2; 3) + (0, 4, 0) \rangle$	$\mathbf{a}, 3\mathbf{b}, \mathbf{c}$	0, 2, 0
[3] $\mathbf{c}' = 3\mathbf{c}$			
$Cmc2_1 (36)$	$\langle (2; 3) + (0, 0, 1) \rangle$	$\mathbf{a}, \mathbf{b}, 3\mathbf{c}$	

• Series of maximal isomorphic subgroups

[p] $\mathbf{a}' = p\mathbf{a}$			
$Cmc2_1 (36)$	$\langle 3; 2 + (2u, 0, 0) \rangle$ prime $p > 2$; $0 \leq u < p$ p conjugate subgroups	$p\mathbf{a}, \mathbf{b}, \mathbf{c}$	$u, 0, 0$
[p] $\mathbf{b}' = p\mathbf{b}$			
$Cmc2_1 (36)$	$\langle (2; 3) + (0, 2u, 0) \rangle$ prime $p > 2$; $0 \leq u < p$ p conjugate subgroups	$\mathbf{a}, p\mathbf{b}, \mathbf{c}$	$0, u, 0$
[p] $\mathbf{c}' = p\mathbf{c}$			
$Cmc2_1 (36)$	$\langle (2; 3) + (0, 0, \frac{p}{2} - \frac{1}{2}) \rangle$ prime $p > 2$ no conjugate subgroups	$\mathbf{a}, \mathbf{b}, p\mathbf{c}$	

I Minimal translationengleiche supergroups

 [2] $Cmcm$ (63); [2] $Cmce$ (64); [3] $P6_3cm$ (185); [3] $P6_3mc$ (186)

II Minimal non-isomorphic klassengleiche supergroups

• Additional centring translations

 [2] $Fmm2$ (42)

• Decreased unit cell

 [2] $\mathbf{a}' = \frac{1}{2}\mathbf{a}$, $\mathbf{b}' = \frac{1}{2}\mathbf{b}$ $Pmc2_1$ (26); [2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $Cmm2$ (35)