

$Pccn$ 

No. 56

 $P2_1/c2_1/c2/n$  $D_{2h}^{10}$ Generators selected (1);  $t(1,0,0)$ ;  $t(0,1,0)$ ;  $t(0,0,1)$ ; (2); (3); (5)

## General position

Multiplicity,  
Wyckoff letter,  
Site symmetry

Coordinates

8  $e$  1 (1)  $x, y, z$  (2)  $\bar{x} + \frac{1}{2}, \bar{y} + \frac{1}{2}, z$  (3)  $\bar{x}, y + \frac{1}{2}, \bar{z} + \frac{1}{2}$  (4)  $x + \frac{1}{2}, \bar{y}, \bar{z} + \frac{1}{2}$   
 (5)  $\bar{x}, \bar{y}, \bar{z}$  (6)  $x + \frac{1}{2}, y + \frac{1}{2}, \bar{z}$  (7)  $x, \bar{y} + \frac{1}{2}, z + \frac{1}{2}$  (8)  $\bar{x} + \frac{1}{2}, y, z + \frac{1}{2}$

I Maximal *translationengleiche* subgroups

[2] $Pc2_1n$ (33, $Pna2_1$ )	1; 3; 6; 8	<b>c, a, b</b>	0, 0, 1/4
[2] $P2_1cn$ (33, $Pna2_1$ )	1; 4; 6; 7	<b>c, b, -a</b>	0, 0, 1/4
[2] $Pcc2$ (27)	1; 2; 7; 8		1/4, 1/4, 0
[2] $P2_12_12$ (18)	1; 2; 3; 4		1/4, 1/4, 1/4
[2] $P12_1/c1$ (14)	1; 3; 5; 7		
[2] $P2_1/c11$ (14, $P12_1/c1$ )	1; 4; 5; 8	<b>-b, a, c</b>	
[2] $P112/n$ (13, $P112/a$ )	1; 2; 5; 6	<b>-a - b, a, c</b>	

II Maximal *klassengleiche* subgroups

## • Enlarged unit cell

[3] $\mathbf{a}' = 3\mathbf{a}$			
$\left\{ \begin{array}{l} Pccn \text{ (56)} \\ Pccn \text{ (56)} \\ Pccn \text{ (56)} \end{array} \right.$	$\langle 3; 5; 2 + (1, 0, 0) \rangle$ $\langle 2 + (3, 0, 0); (3; 5) + (2, 0, 0) \rangle$ $\langle 2 + (5, 0, 0); (3; 5) + (4, 0, 0) \rangle$	<b>3a, b, c</b> <b>3a, b, c</b> <b>3a, b, c</b>	1, 0, 0 2, 0, 0
[3] $\mathbf{b}' = 3\mathbf{b}$			
$\left\{ \begin{array}{l} Pccn \text{ (56)} \\ Pccn \text{ (56)} \\ Pccn \text{ (56)} \end{array} \right.$	$\langle 5; (2; 3) + (0, 1, 0) \rangle$ $\langle 2 + (0, 3, 0); 3 + (0, 1, 0); 5 + (0, 2, 0) \rangle$ $\langle 2 + (0, 5, 0); 3 + (0, 1, 0); 5 + (0, 4, 0) \rangle$	<b>a, 3b, c</b> <b>a, 3b, c</b> <b>a, 3b, c</b>	0, 1, 0 0, 2, 0
[3] $\mathbf{c}' = 3\mathbf{c}$			
$\left\{ \begin{array}{l} Pccn \text{ (56)} \\ Pccn \text{ (56)} \\ Pccn \text{ (56)} \end{array} \right.$	$\langle 2; 5; 3 + (0, 0, 1) \rangle$ $\langle 2; 3 + (0, 0, 3); 5 + (0, 0, 2) \rangle$ $\langle 2; 3 + (0, 0, 5); 5 + (0, 0, 4) \rangle$	<b>a, b, 3c</b> <b>a, b, 3c</b> <b>a, b, 3c</b>	0, 0, 1 0, 0, 2

## • Series of maximal isomorphic subgroups

[ $p$ ] $\mathbf{a}' = p\mathbf{a}$			
$Pccn$ (56)	$\langle 2 + (0, \frac{p}{2} - \frac{1}{2} + 2u, 0, 0); (3; 5) + (2u, 0, 0) \rangle$ prime $p > 2$ ; $0 \leq u < p$ $p$ conjugate subgroups	<b><math>pa, b, c</math></b>	$u, 0, 0$
[ $p$ ] $\mathbf{b}' = p\mathbf{b}$			
$Pccn$ (56)	$\langle 2 + (0, \frac{p}{2} - \frac{1}{2} + 2u, 0); 3 + (0, \frac{p}{2} - \frac{1}{2}, 0); 5 + (0, 2u, 0) \rangle$ prime $p > 2$ ; $0 \leq u < p$ $p$ conjugate subgroups	<b><math>a, pb, c</math></b>	$0, u, 0$
[ $p$ ] $\mathbf{c}' = p\mathbf{c}$			
$Pccn$ (56)	$\langle 2; 3 + (0, 0, \frac{p}{2} - \frac{1}{2} + 2u); 5 + (0, 0, 2u) \rangle$ prime $p > 2$ ; $0 \leq u < p$ $p$ conjugate subgroups	<b><math>a, b, pc</math></b>	$0, 0, u$

I Minimal *translationengleiche* supergroups[2]  $P4/ncc$  (130); [2]  $P4_2/ncm$  (138)II Minimal non-isomorphic *klassengleiche* supergroups

## • Additional centring translations

[2]  $Aema$  (64,  $Cmce$ ); [2]  $Bmeb$  (64,  $Cmce$ ); [2]  $Cccm$  (66); [2]  $Ibam$  (72)

## • Decreased unit cell

[2]  $\mathbf{a}' = \frac{1}{2}\mathbf{a}$   $Pccb$  (54,  $Pcca$ ); [2]  $\mathbf{b}' = \frac{1}{2}\mathbf{b}$   $Pcca$  (54); [2]  $\mathbf{c}' = \frac{1}{2}\mathbf{c}$   $Pmmn$  (59)