

S_4^2 $I\bar{4}$

No. 82

 $I\bar{4}$ Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; $t(\frac{1}{2},\frac{1}{2},\frac{1}{2})$; (2); (3)

General position

Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

 $(0,0,0)+$ $(\frac{1}{2},\frac{1}{2},\frac{1}{2})+$ 8 g 1(1) x,y,z (2) \bar{x},\bar{y},z (3) y,\bar{x},\bar{z} (4) \bar{y},x,\bar{z} I Maximal *translationengleiche* subgroups[2] $I2$ (5, A112)

(1; 2)+

 $\mathbf{b}, -\mathbf{a} - \mathbf{b}, \mathbf{c}$ II Maximal *klassengleiche* subgroups

• Loss of centring translations

[2] $P\bar{4}$ (81)

1; 2; 3; 4

[2] $P\bar{4}$ (81)1; 2; (3; 4) + $(\frac{1}{2},\frac{1}{2},\frac{1}{2})$

1/2, 0, 1/4

• Enlarged unit cell

[3] $\mathbf{c}' = 3\mathbf{c}$ $I\bar{4}$ (82) $\langle 2; 3 \rangle$ $\mathbf{a}, \mathbf{b}, 3\mathbf{c}$ $I\bar{4}$ (82) $\langle 2; 3 + (0,0,2) \rangle$ $\mathbf{a}, \mathbf{b}, 3\mathbf{c}$

0, 0, 1

 $I\bar{4}$ (82) $\langle 2; 3 + (0,0,4) \rangle$ $\mathbf{a}, \mathbf{b}, 3\mathbf{c}$

0, 0, 2

• Series of maximal isomorphic subgroups

[p] $\mathbf{c}' = p\mathbf{c}$ $I\bar{4}$ (82) $\langle 2; 3 + (0,0,2u) \rangle$ $\mathbf{a}, \mathbf{b}, p\mathbf{c}$ 0, 0, u prime $p > 2$; $0 \leq u < p$ p conjugate subgroups[p^2] $\mathbf{a}' = p\mathbf{a}, \mathbf{b}' = p\mathbf{b}$ $I\bar{4}$ (82) $\langle 2 + (2u, 2v, 0); 3 + (u - v, u + v, 0) \rangle$ $p\mathbf{a}, p\mathbf{b}, \mathbf{c}$ $u, v, 0$ prime $p > 2$; $0 \leq u < p$; $0 \leq v < p$ p^2 conjugate subgroups for $p = 4n - 1$ [$p = q^2 + r^2$] $\mathbf{a}' = q\mathbf{a} - r\mathbf{b}, \mathbf{b}' = r\mathbf{a} + q\mathbf{b}$ $I\bar{4}$ (82) $\langle 2 + (2u, 0, 0); 3 + (u, u, 0) \rangle$ $q\mathbf{a} - r\mathbf{b}, r\mathbf{a} + q\mathbf{b}, \mathbf{c}$ $u, 0, 0$ prime $p > 4$; $q > 0$; $r > 0$; $0 \leq u < p$ p conjugate subgroups for $p = 4n + 1$ I Minimal *translationengleiche* supergroups[2] $I4/m$ (87); [2] $I4_1/a$ (88); [2] $I\bar{4}m2$ (119); [2] $I\bar{4}c2$ (120); [2] $I\bar{4}2m$ (121); [2] $I\bar{4}2d$ (122)II Minimal non-isomorphic *klassengleiche* supergroups

• Additional centring translations

none

• Decreased unit cell

[2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $C\bar{4}$ (81, $P\bar{4}$)