

D_{4h}^{14}
 $P4_2/m2_1/n2/m$

No. 136

 $P4_2/mnm$
Generators selected (1); $t(1,0,0)$; $t(0,1,0)$; $t(0,0,1)$; (2); (3); (5); (9)

General position

 Multiplicity,
Wyckoff letter,
Site symmetry

Coordinates

16	<i>k</i>	1	(1) x, y, z	(2) \bar{x}, \bar{y}, z	(3) $\bar{y} + \frac{1}{2}, x + \frac{1}{2}, z + \frac{1}{2}$	(4) $y + \frac{1}{2}, \bar{x} + \frac{1}{2}, z + \frac{1}{2}$
			(5) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, \bar{z} + \frac{1}{2}$	(6) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, \bar{z} + \frac{1}{2}$	(7) y, x, \bar{z}	(8) $\bar{y}, \bar{x}, \bar{z}$
			(9) $\bar{x}, \bar{y}, \bar{z}$	(10) x, y, \bar{z}	(11) $y + \frac{1}{2}, \bar{x} + \frac{1}{2}, \bar{z} + \frac{1}{2}$	(12) $\bar{y} + \frac{1}{2}, x + \frac{1}{2}, \bar{z} + \frac{1}{2}$
			(13) $x + \frac{1}{2}, \bar{y} + \frac{1}{2}, z + \frac{1}{2}$	(14) $\bar{x} + \frac{1}{2}, y + \frac{1}{2}, z + \frac{1}{2}$	(15) \bar{y}, \bar{x}, z	(16) y, x, z

I Maximal translationengleiche subgroups

[2] $P\bar{4}n2$ (118)	1; 2; 7; 8; 11; 12; 13; 14		0, 1/2, 1/4
[2] $P4_2m$ (113)	1; 2; 5; 6; 11; 12; 15; 16		0, 1/2, 1/4
[2] $P4_2nm$ (102)	1; 2; 3; 4; 13; 14; 15; 16		
[2] $P4_22_12$ (94)	1; 2; 3; 4; 5; 6; 7; 8		
[2] $P4_2/m11$ (84, $P4_2/m$)	1; 2; 3; 4; 9; 10; 11; 12		0, 1/2, 0
[2] $P2/m12/m$ (65, $\bar{C}mmm$)	1; 2; 7; 8; 9; 10; 15; 16	a - b, a + b, c	
[2] $P2/m2_1/n1$ (58, $Pnmm$)	1; 2; 5; 6; 9; 10; 13; 14		

II Maximal klassengleiche subgroups

• Enlarged unit cell

[3] $\mathbf{c}' = 3\mathbf{c}$			
$\left\{ \begin{array}{l} P4_2/mnm \text{ (136)} \\ P4_2/mnm \text{ (136)} \\ P4_2/mnm \text{ (136)} \end{array} \right.$	$\langle 2; 9; (3; 5) + (0, 0, 1) \rangle$	a, b, 3c	
	$\langle 2; 3 + (0, 0, 1); 5 + (0, 0, 3); 9 + (0, 0, 2) \rangle$	a, b, 3c	0, 0, 1
	$\langle 2; 3 + (0, 0, 1); 5 + (0, 0, 5); 9 + (0, 0, 4) \rangle$	a, b, 3c	0, 0, 2

• Series of maximal isomorphic subgroups

[<i>p</i>] $\mathbf{c}' = p\mathbf{c}$			
$P4_2/mnm$ (136)	$\langle 2; 3 + (0, 0, \frac{p}{2} - \frac{1}{2}); 5 + (0, 0, \frac{p}{2} - \frac{1}{2} + 2u); 9 + (0, 0, 2u) \rangle$	a, b, pc	0, 0, <i>u</i>
	prime $p > 2$; $0 \leq u < p$ <i>p</i> conjugate subgroups		
[p^2] $\mathbf{a}' = p\mathbf{a}, \mathbf{b}' = p\mathbf{b}$			
$P4_2/mnm$ (136)	$\langle (2; 9) + (2u, 2v, 0); 3 + (\frac{p}{2} - \frac{1}{2} + u + v, \frac{p}{2} - \frac{1}{2} - u + v, 0); 5 + (\frac{p}{2} - \frac{1}{2} + 2u, \frac{p}{2} - \frac{1}{2}, 0) \rangle$	pa, pb, c	<i>u, v, 0</i>
	prime $p > 2$; $0 \leq u < p$; $0 \leq v < p$		
	p^2 conjugate subgroups		

I Minimal translationengleiche supergroups

none

II Minimal non-isomorphic klassengleiche supergroups

• Additional centring translations

 [2] $C4_2/mcm$ (131, $P4_2/mmc$); [2] $I4/mmm$ (139)

• Decreased unit cell

 [2] $\mathbf{c}' = \frac{1}{2}\mathbf{c}$ $P4/mbm$ (127)