


1.4. SPACE GROUPS AND THEIR DESCRIPTIONS



**Figure 1.4.4.3** Symmetry-element diagram for the space group  $P4$  (75) for the orthogonal projection along  $[001]$ .

can not contain  $Y = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{pmatrix}$  and thus  $S_X$  and  $S_Y$  are not conjugate in  $\mathcal{G}$ .

However, the translation by  $(\frac{1}{2}, \frac{1}{2}, 0)$  conjugates  $S_X$  to  $S_Y$ , while fixing the group  $\mathcal{G}$  as a whole. This shows that there is an ambiguity in choosing the origin either at  $0, 0, 0$  or  $\frac{1}{2}, \frac{1}{2}, 0$ , since these points are geometrically indistinguishable (both being intersections of a fourfold axis with the  $ab$  plane).

The ambiguity in the origin choice in the above example can be explained by the *affine normalizer* of the space group  $\mathcal{G}$  (see Section 1.1.8 for a general introduction to normalizers). The full group  $\mathcal{A}$  of affine mappings acts *via* conjugation on the set of space groups and the space groups of the same affine type are obtained as the orbit of a single group of that type under  $\mathcal{A}$ .

*Definition*

The group  $\mathcal{N}$  of affine mappings  $n \in \mathcal{A}$  that fix a space group  $\mathcal{G}$  under conjugation is called the *affine normalizer* of  $\mathcal{G}$ , *i.e.*

$$\mathcal{N} = \mathcal{N}_{\mathcal{A}}(\mathcal{G}) = \{n \in \mathcal{A} | n\mathcal{G}n^{-1} = \mathcal{G}\}.$$

The affine normalizer is the largest subgroup of  $\mathcal{A}$  such that  $\mathcal{G}$  is a normal subgroup of  $\mathcal{N}$ .

Conjugation by operations of the affine normalizer results in a permutation of the operations of  $\mathcal{G}$ , *i.e.* in a relabelling without changing their geometric properties. The additional translations contained in the affine normalizer can in fact be derived from the space-group diagrams, because shifting the origin by such a translation results in precisely the same diagram. More generally, an element of the affine normalizer can be interpreted as a change of the coordinate system that does not alter the space-group diagrams.

A more thorough description of the affine normalizers of space groups is given in Chapter 3.5, where tables with the affine normalizers are also provided.

Since the affine normalizer of a space group  $\mathcal{G}$  is in general a group containing  $\mathcal{G}$  as a proper subgroup, it is possible that subgroups of  $\mathcal{G}$  that are not conjugate by any operation of  $\mathcal{G}$  may be conjugate by an operation in the affine normalizer. As a consequence, the site-symmetry groups  $S_X$  and  $S_Y$  of two points  $X$  and  $Y$  belonging to different Wyckoff positions of  $\mathcal{G}$  may be conjugate under the affine normalizer of  $\mathcal{G}$ . This reveals that the points  $X$  and  $Y$  are in fact geometrically equivalent, since they fall into the same orbit under the affine normalizer of  $\mathcal{G}$ . Joining the equivalence classes of these points into a single class results in a coarser classification with larger classes, which are called *Wyckoff sets*.

*Definition*

Two points  $X$  and  $Y$  belong to the same *Wyckoff set* if their site-symmetry groups  $S_X$  and  $S_Y$  are conjugate subgroups of the affine normalizer  $\mathcal{N}_{\mathcal{A}}(\mathcal{G})$  of  $\mathcal{G}$ .


In particular, the Wyckoff set containing a point  $X$  also contains the full orbit of  $X$  under the affine normalizer of  $\mathcal{G}$ .

*Example*

Let  $\mathcal{G}$  be the space group of type  $P222_1$  (17) generated by the translations of an orthorhombic lattice, the twofold rotation  $\{2_{100}|0\}: x, \bar{y}, \bar{z}$  and the twofold screw rotation  $\{2_{001}|0, 0, \frac{1}{2}\}: \bar{x}, \bar{y}, z + \frac{1}{2}$ . Note that the composition of these two elements is the twofold rotation with the line  $0, y, \frac{1}{4}$  as its geometric element. The group  $\mathcal{G}$  has four different Wyckoff positions with a site-symmetry group generated by a twofold rotation; representatives of these Wyckoff positions are the

points  $X_1 = \begin{pmatrix} x \\ 0 \\ 0 \end{pmatrix}$  (Wyckoff position  $2a$ , site-symmetry symbol  $2..$ ),  $X_2 = \begin{pmatrix} x \\ \frac{1}{2} \\ 0 \end{pmatrix}$  (position  $2b$ , symbol  $2..$ ),  $Y_1 = \begin{pmatrix} 0 \\ y \\ \frac{1}{4} \end{pmatrix}$  (position  $2c$ , symbol  $.2.$ ) and  $Y_2 = \begin{pmatrix} 0 \\ y \\ \frac{1}{4} \end{pmatrix}$  (position  $2d$ , symbol  $.2.$ ).

From the tables of affine normalizers in Chapter 3.5, but also by a careful analysis of the space-group diagrams in Fig. 1.4.4.4, one deduces that the affine normalizer of  $\mathcal{G}$  contains the additional translations  $t(\frac{1}{2}, 0, 0)$ ,  $t(0, \frac{1}{2}, 0)$  and  $t(0, 0, \frac{1}{2})$ , since all the diagrams are invariant by a shift of  $\frac{1}{2}$  along any of the coordinate axes. Moreover, the symmetry operation  $\{m_{\bar{1}\bar{1}0}|0, 0, \frac{1}{4}\}: y, x, z + \frac{1}{4}$  which interchanges the  $a$  and  $b$  axes and shifts the origin by  $\frac{1}{4}$  along the  $c$  axis belongs to the affine



**Figure 1.4.4.4** Symmetry-element diagrams for the space group  $P222_1$  (17) for orthogonal projections along  $[001]$ ,  $[010]$ ,  $[100]$  (left to right).