

1.4. SPACE GROUPS AND THEIR DESCRIPTIONS

some practice each of the diagrams can be generated from the other. In a number of texts, the two diagrams are considered as completely equivalent descriptions of the same space group. This statement is true for most of the space groups. However, there are a number of space groups for which the point configuration displayed on the general-position diagram has higher symmetry than the generating space group (Suescun & Nespolo, 2012; Müller, 2012). For example, consider the diagrams of the space group $P2$, No. 3 (unique axis b , cell choice 1) shown in Fig. 1.4.2.7. It is easy to recognise that, apart from the twofold rotations, the point configuration shown in the general-position diagram is symmetric with respect to a reflection through a plane containing the general-position points, and as a result the space group of the general-position configuration is of $P2/m$ type, and not of $P2$. There are a number of space groups for which the general-position diagram displays higher space-group symmetry, for example: $P1$, $P2_1$, $P4mm$, $P6$ etc. The analysis of the eigensymmetry groups of the general-position orbits results in a systematic procedure for the determination of such space groups: the general-position diagrams do not reflect the space-group symmetry correctly if the general-position orbits are *non-characteristic*, i.e. their eigensymmetry groups are supergroups of the space groups. (An introduction to terms like eigensymmetry groups, characteristic and non-characteristic orbits, and further discussion of space groups with non-characteristic general-position orbits are given in Section 1.4.4.4.)

- (2) The graphical presentation of the general-position points of cubic groups is more difficult: three different parameters are required to specify the height of the points along the projection direction. To make the presentation clearer, the general-position points are grouped around points of higher site symmetry and represented in the form of polyhedra. For most of the space groups the initial general point is taken as 0.048, 0.12, 0.089, and the polyhedra are centred at 0, 0, 0 (and its equivalent points). Additional general-position diagrams are shown for space groups with special sites different from 0, 0, 0 that have site-symmetry groups of equal or higher order. Consider, for example, the two general-position diagrams of the space group $I4_132$ (214) shown in Fig. 1.4.2.8. The polyhedra of the left-hand diagram are centred at special points of highest site-symmetry, namely, at $\frac{1}{8}, \frac{1}{8}, \frac{1}{8}$ and its equivalent points in the unit cell. The site-symmetry groups are of the type 32 leading to polyhedra in the form of *twisted trigonal antiprisms* (cf. Table 3.2.3.2). The polyhedra (sphenoids) of the right-hand diagram are attached to the origin 0, 0, 0 and its equivalent points in the unit cell, site-symmetry group of the type 3. The fractions attached to the polyhedra indicate the heights of the high-symmetry points along the projection direction (cf. Section 2.1.3.6 for further explanations of the diagrams).

1.4.3. Generation of space groups

BY H. WONDRATSCHKE

In group theory, a *set of generators* of a group is a set of group elements such that each group element may be obtained as a finite ordered product of the generators. For space groups of one, two and three dimensions, generators may always be chosen and

Table 1.4.3.1

Sequence of generators for the crystal classes

The space-group generators differ from those listed here by their glide or screw components. The generator 1 is omitted, except for crystal class 1. The generators are represented by the corresponding Seitz symbols (cf. Tables 1.4.2.1–1.4.2.3). Following the conventions, the subscript of a symbol denotes the characteristic direction of that operation, where necessary. For example, the subscripts 001, 010, 110 etc. refer to the directions [001], [010], [110] etc. For mirror reflections m , the ‘direction of m ’ refers to the normal of the mirror plane.

Hermann–Mauguin symbol of crystal class	Generators g_i (sequence left to right)
1 $\bar{1}$	1 $\bar{1}$
2 m $2/m$	2 m 2, $\bar{1}$
222 $mm2$ mmm	$2_{001}, 2_{010}$ $2_{001}, m_{010}$ $2_{001}, 2_{010}, \bar{1}$
4 $\bar{4}$ $4/m$ 422 $4mm$ $\bar{4}2m$ $\bar{4}m2$ $4/mmm$	$2_{001}, 4_{001}^+$ $2_{001}, 4_{001}^+$ $2_{001}, 4_{001}^+, \bar{1}$ $2_{001}, 4_{001}^+, 2_{010}$ $2_{001}, 4_{001}^+, m_{010}$ $2_{001}, 4_{001}^+, 2_{010}$ $2_{001}, 4_{001}^+, m_{010}$ $2_{001}, 4_{001}^+, 2_{010}, \bar{1}$
3 (rhombohedral coordinates) $\bar{3}$ (rhombohedral coordinates) 321 (rhombohedral coordinates) 312 $3m1$ (rhombohedral coordinates) $31m$ $\bar{3}m1$ (rhombohedral coordinates) $\bar{3}1m$	3_{001}^+ 3_{111}^+ $3_{001}^+, \bar{1}$ $3_{111}^+, \bar{1}$ $3_{001}^+, 2_{110}$ $3_{111}^+, 2_{101}$ $3_{001}^+, 2_{1\bar{1}0}$ $3_{001}^+, m_{110}$ $3_{111}^+, m_{101}$ $3_{001}^+, m_{1\bar{1}0}$ $3_{001}^+, 2_{110}, \bar{1}$ $3_{111}^+, 2_{101}, \bar{1}$ $3_{001}^+, 2_{1\bar{1}0}, \bar{1}$
6 $\bar{6}$ $6/m$ 622 $6mm$ $\bar{6}m2$ $\bar{6}2m$ $6/mmm$	$3_{001}^+, 2_{001}$ $3_{001}^+, m_{001}$ $3_{001}^+, 2_{001}, \bar{1}$ $3_{001}^+, 2_{001}, 2_{110}$ $3_{001}^+, 2_{001}, m_{110}$ $3_{001}^+, m_{001}, m_{110}$ $3_{001}^+, m_{001}, 2_{110}$ $3_{001}^+, 2_{001}, 2_{110}, \bar{1}$
23 $m\bar{3}$ 432 $\bar{4}3m$ $m\bar{3}m$	$2_{001}, 2_{010}, 3_{111}^+$ $2_{001}, 2_{010}, 3_{111}^+, \bar{1}$ $2_{001}, 2_{010}, 3_{111}^+, 2_{110}$ $2_{001}, 2_{010}, 3_{111}^+, m_{1\bar{1}0}$ $2_{001}, 2_{010}, 3_{111}^+, 2_{110}, \bar{1}$

ordered in such a way that each symmetry operation W can be written as the product of powers of h generators g_j ($j = 1, 2, \dots, h$). Thus,

$$W = g_h^{k_h} \cdot g_{h-1}^{k_{h-1}} \cdot \dots \cdot g_p^{k_p} \cdot \dots \cdot g_3^{k_3} \cdot g_2^{k_2} \cdot g_1,$$

where the powers k_j are positive or negative integers (including zero). The description of a group by means of generators has the advantage of compactness. For instance, the 48 symmetry operations in point group $m\bar{3}m$ can be described by two

1. INTRODUCTION TO SPACE-GROUP SYMMETRY

Table 1.4.3.2

Generation of the space group $P6_122 \equiv D_6^2$ (178)

The entries in the second column designated by the numbers (1)–(12) correspond to the coordinate triplets of the general position of $P6_122$.

	Coordinate triplets	Symmetry operations
g_1	(1) x, y, z ;	Identity I
g_2	$\left. \begin{array}{l} t(100) \\ t(010) \\ t(001) \end{array} \right\}$ The group $\mathcal{G}_4 \equiv T$ of all translations of $P6_122$ has been generated	$\left\{ \begin{array}{l} \\ \\ \\ \end{array} \right.$ Generating translations
g_3		
g_4		
g_5	(2) $\bar{y}, x - y, z + \frac{1}{3}$;	Threefold screw rotation
g_5^2	(3) $\bar{x} + y, \bar{x}, z + \frac{2}{3}$;	Threefold screw rotation
$g_5^3 = t(001)$	Now the space group $\mathcal{G}_5 \equiv P3_1$ has been generated	
g_6	(4) $\bar{x}, \bar{y}, z + \frac{1}{2}$;	Twofold screw rotation
$g_6 * g_5$	(5) $y, \bar{x} + y, z + \frac{5}{6}$;	Sixfold screw rotation
$g_6 * g_5^2$	$x - y, x, z + \frac{7}{6} \sim$ (6) $x - y, x, z + \frac{1}{6}$;	Sixfold screw rotation
$g_6^2 = t(001)$	Now the space group $\mathcal{G}_6 \equiv P6_1$ has been generated	
g_7	(7) $y, x, \bar{z} + \frac{1}{3}$;	Twofold rotation, direction of axis [110]
$g_7 * g_5$	(8) $x - y, \bar{y}, \bar{z}$;	Twofold rotation, axis [100]
$g_7 * g_5^2$	$\bar{x}, \bar{x} + y, \bar{z} - \frac{1}{3} \sim$ (9) $\bar{x}, \bar{x} + y, \bar{z} + \frac{2}{3}$;	Twofold rotation, axis [010]
$g_7 * g_6$	$\bar{y}, \bar{x}, \bar{z} - \frac{1}{6} \sim$ (10) $\bar{y}, \bar{x}, \bar{z} + \frac{5}{6}$;	Twofold rotation, axis [110]
$g_7 * g_6 * g_5$	$\bar{x} + y, y, \bar{z} - \frac{1}{2} \sim$ (11) $\bar{x} + y, y, \bar{z} + \frac{1}{2}$;	Twofold rotation, axis [120]
$g_7 * g_6 * g_5^2$	$x, x - y, \bar{z} - \frac{5}{6} \sim$ (12) $x, x - y, \bar{z} + \frac{1}{6}$;	Twofold rotation, axis [210]
$g_7^2 = I$	$\mathcal{G}_7 \sim P6_122$	

generators. Different choices of generators are possible. For the space-group tables, generators and generating procedures have been chosen such as to make the entries in the blocks ‘General position’ (cf. Section 2.1.3.11) and ‘Symmetry operations’ (cf. Section 2.1.3.9) as transparent as possible. Space groups of the same crystal class are generated in the same way (see Table 1.4.3.1 for the sequences that have been chosen), and the aim has been to accentuate important subgroups of space groups as much as possible. Accordingly, a process of generation in the form of a *composition series* has been adopted, see Ledermann (1976). The generator g_1 is defined as the identity operation, represented by (1) x, y, z . The generators g_2, g_3 , and g_4 are the translations with translation vectors \mathbf{a}, \mathbf{b} and \mathbf{c} , respectively. Thus, the coefficients k_2, k_3 and k_4 may have any integral value. If centring translations exist, they are generated by translations g_5 (and g_6 in the case of an F lattice) with translation vectors \mathbf{d} (and \mathbf{e}). For a C lattice, for example, \mathbf{d} is given by $\mathbf{d} = \frac{1}{2}(\mathbf{a} + \mathbf{b})$. The exponents k_5 (and k_6) are restricted to the following values:

Lattice letter A, B, C, I : $k_5 = 0$ or 1 .

Lattice letter R (hexagonal axes): $k_5 = 0, 1$ or 2 .

Lattice letter F : $k_5 = 0$ or 1 ; $k_6 = 0$ or 1 .

As a consequence, any translation t of \mathcal{G} with translation vector

$$\mathbf{t} = k_2\mathbf{a} + k_3\mathbf{b} + k_4\mathbf{c} (+ k_5\mathbf{d} + k_6\mathbf{e})$$

can be obtained as a product

$$t = (g_6)^{k_6} \cdot (g_5)^{k_5} \cdot g_4^{k_4} \cdot g_3^{k_3} \cdot g_2^{k_2} \cdot g_1,$$

where k_2, \dots, k_6 are integers determined by t . The generators g_6 and g_5 are enclosed between parentheses because they are effective only in centred lattices.

The remaining generators generate those symmetry operations that are not translations. They are chosen in such a way that only terms g_j or g_j^2 occur. For further specific rules, see below.

The process of generating the entries of the space-group tables may be demonstrated by the example in Table 1.4.3.2, where \mathcal{G}_j denotes the group generated by g_1, g_2, \dots, g_j . For $j \geq 5$, the next generator g_{j+1} is introduced when $g_j^{k_j} \in \mathcal{G}_{j-1}$, because

in this case no new symmetry operation would be generated by $g_j^{k_j}$. The generating process is terminated when there is no further generator. In the present example, g_7 completes the generation: $\mathcal{G}_7 \equiv P6_122$ (178).

1.4.3.1. Selected order for non-translational generators

For the non-translational generators, the following sequence has been adopted:

- (a) In all centrosymmetric space groups, an inversion (if possible at the origin O) has been selected as the last generator.
- (b) Rotations precede symmetry operations of the second kind. In crystal classes $\bar{4}2m$ and $4m2$ and $\bar{6}2m$ and $\bar{6}m2$, as an exception, $\bar{4}$ and $\bar{6}$ are generated first in order to take into account the conventional choice of origin in the fixed points of $\bar{4}$ and $\bar{6}$.
- (c) The non-translational generators of space groups with C, A, B, F, I or R symbols are those of the corresponding space group with a P symbol, if possible. For instance, the generators of $I2_12_12_1$ (24) are those of $P2_12_12_1$ (19) and the generators of $Ibca$ (73) are those of $Pbca$ (61), apart from the centring translations.

Exceptions: $I4cm$ (108) and $I4/mcm$ (140) are generated via $P4cc$ (103) and $P4/mcc$ (124), because $P4cm$ and $P4/mcm$ do not exist. In space groups with d glides (except $I\bar{4}2d$, No. 122) and also in $I4_1/a$ (88), the corresponding rotation subgroup has been generated first. The generators of this subgroup are the same as those of the corresponding space group with a lattice symbol P .

Example

$F4_1/d\bar{3}2/m$ (227):

$P4_132$ (213) \longrightarrow $F4_132$ (210) \longrightarrow $F4_1/d\bar{3}2/m$.

- (d) In some cases, rule (c) could not be followed without breaking rule (a), e.g. in $Cmme$ (67). In such cases, the generators are chosen to correspond to the Hermann–Mauguin symbol as far as possible. For instance, the generators (apart from centring) of $Cmme$ and $Imma$ (74) are

those of $Pmmb$, which is a non-standard setting of $Pmma$ (51). (A combination of the generators of $Pmma$ with the C - or I -centring translation results in non-standard settings of $Cmme$ and $Imma$.)

For the space groups with lattice symbol P , the generation procedure has given the same triplets (except for their sequence) as in *IT* (1952). In non- P space groups, the triplets listed sometimes differ from those of *IT* (1952) by a centring translation.

1.4.4. General and special Wyckoff positions

BY B. SOUVIGNIER

One of the first tasks in the analysis of crystal patterns is to determine the actual positions of the atoms. Since the full crystal pattern can be reconstructed from a single unit cell or even an asymmetric unit, it is clearly sufficient to focus on the atoms inside such a restricted volume. What one observes is that the atoms typically do not occupy arbitrary positions in the unit cell, but that they often lie on geometric elements, *e.g.* reflection planes or lines along rotation axes. It is therefore very useful to analyse the symmetry properties of the points in a unit cell in order to predict likely positions of atoms.

We note that in this chapter all statements and definitions refer to the usual three-dimensional space \mathbb{E}^3 , but also can be formulated, *mutatis mutandis*, for plane groups acting on \mathbb{E}^2 and for higher-dimensional groups acting on n -dimensional space \mathbb{E}^n .

1.4.4.1. Crystallographic orbits

Since the operations of a space group provide symmetries of a crystal pattern, two points X and Y that are mapped onto each other by a space-group operation are regarded as being *geometrically equivalent*. Starting from a point $X \in \mathbb{E}^3$, infinitely many points Y equivalent to X are obtained by applying all space-group operations $g = (\mathbf{W}, \mathbf{w})$ to X : $Y = g(X) = (\mathbf{W}, \mathbf{w})X = (\mathbf{W}X + \mathbf{w})$.

Definition

For a space group \mathcal{G} acting on the three-dimensional space \mathbb{E}^3 , the (infinite) set

$$\mathcal{O} = \mathcal{G}(X) := \{g(X) | g \in \mathcal{G}\}$$

is called the *orbit of X under \mathcal{G}* .

The orbit of X is the smallest subset of \mathbb{E}^3 that contains X and is closed under the action of \mathcal{G} . It is also called a *crystallographic orbit*.

Every point in direct space \mathbb{E}^3 belongs to precisely one orbit under \mathcal{G} and thus the orbits of \mathcal{G} partition the direct space into disjoint subsets. It is clear that an orbit is completely determined by its points in the unit cell, since translating the unit cell by the translation subgroup \mathcal{T} of \mathcal{G} entirely covers \mathbb{E}^3 .

It may happen that two different symmetry operations g and h in \mathcal{G} map X to the same point. Since $g(X) = h(X)$ implies that $h^{-1}g(X) = X$, the point X is fixed by the nontrivial operation $h^{-1}g$ in \mathcal{G} .

Definition

The subgroup $\mathcal{S}_X = \mathcal{S}_{\mathcal{G}}(X) := \{g \in \mathcal{G} | g(X) = X\}$ of symmetry operations from \mathcal{G} that fix X is called the *site-symmetry group of X in \mathcal{G}* .

Since translations, glide reflections and screw rotations fix no point in \mathbb{E}^3 , a site-symmetry group \mathcal{S}_X never contains operations of these types and thus consists only of reflections, rotations, inversions and rotoinversions. Because of the absence of translations, \mathcal{S}_X contains at most one operation from a coset $\mathcal{T}g$ relative to the translation subgroup \mathcal{T} of \mathcal{G} , since otherwise the quotient of two such operations tg and $t'g$ would be the non-trivial translation $tgg^{-1}t'^{-1} = tt'^{-1}$ (see Chapter 1.3 for a discussion of coset decompositions). In particular, the operations in \mathcal{S}_X all have different linear parts and because these linear parts form a subgroup of the point group \mathcal{P} of \mathcal{G} , the order of the site-symmetry group \mathcal{S}_X is a divisor of the order of the point group of \mathcal{G} .

The site-symmetry group of a point X is thus a finite subgroup of the space group \mathcal{G} , a subgroup which is isomorphic to a subgroup of the point group \mathcal{P} of \mathcal{G} .

Example

For a space group \mathcal{G} of type $P\bar{1}$, the site-symmetry group of the

origin $X = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ is clearly generated by the inversion in the origin: $\{\bar{1}|0\}(X) = X$. On the other hand, the point $Y = \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \end{pmatrix}$

is fixed by the inversion in Y , *i.e.*

$$\{\bar{1}|1, 0, 1\}(Y) = \begin{pmatrix} \bar{1} & 0 & 0 \\ 0 & \bar{1} & 0 \\ 0 & 0 & \bar{1} \end{pmatrix} \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \end{pmatrix} = Y.$$

The symmetry operation $\{\bar{1}|1, 0, 1\}$ also belongs to \mathcal{G} and generates the site-symmetry group of Y . The site-symmetry groups $\mathcal{S}_X = \{\{1|0\}, \{\bar{1}|0\}\}$ of X and $\mathcal{S}_Y = \{\{1|0\}, \{\bar{1}|1, 0, 1\}\}$ of Y are thus different subgroups of order 2 of \mathcal{G} which are isomorphic to the point group of \mathcal{G} (which is generated by $\bar{1}$).

The order $|\mathcal{S}_X|$ of the site-symmetry group \mathcal{S}_X is closely related to the number of points in the orbit of X that lie in the unit cell. An application of the orbit-stabilizer theorem (see Section 1.1.7) yields the crucial observation that each point $Y = g(X)$ in the orbit of X under \mathcal{G} is obtained precisely $|\mathcal{S}_X|$ times as an orbit point: for each $h \in \mathcal{S}_X$ one has $gh(X) = g(X) = Y$ and conversely $g'(X) = g(X)$ implies that $g^{-1}g' = h \in \mathcal{S}_X$ and thus $g' = gh$ for an operation h in \mathcal{S}_X .

Assuming first that we are dealing with a space group \mathcal{G} described by a *primitive* lattice, each coset of \mathcal{G} relative to the translation subgroup \mathcal{T} contains precisely one operation g such that $g(X)$ lies in the primitive unit cell. Since the number of cosets equals the order $|\mathcal{P}|$ of the point group \mathcal{P} of \mathcal{G} and since each orbit point is obtained $|\mathcal{S}_X|$ times, it follows that the number of orbit points in the unit cell is $|\mathcal{P}|/|\mathcal{S}_X|$.

If we deal with a space group with a centred unit cell, the above result has to be modified slightly. If there are $k - 1$ centring vectors, the lattice spanned by the conventional basis is a sublattice of index k in the full translation lattice. The conventional cell therefore is built up from k primitive unit cells (spanned by a primitive lattice basis) and thus in particular contains k times as many points as the primitive cell (see Chapter 1.3 for a detailed discussion of conventional and primitive bases and cells).