

1.5. TRANSFORMATIONS OF COORDINATE SYSTEMS

mined lattice parameters of the low-symmetry phase [$a_{\text{hex}} = 4.164$ (2) Å, $c_{\text{hex}} = 10.69$ (4) Å (Chattopadhyay *et al.*, 1987)] determines the lattice deformation accompanying the displacive phase transition, which basically consists of expanding the cubic unit cell along the [111] direction. (In fact, the elongation along [111] is accompanied by a contraction in the ab plane that leads to an overall volume reduction of about 1.3%.)

Owing to the polar character of $R3m$, the symmetry conditions following from the group–subgroup relation $Fm\bar{3}m > R3m$ [cf. equation (1.5.2.11)] are not sufficient to determine the origin shift of the transformation between the high- and the low-symmetry space groups. The origin shift of $\mathbf{p} = (-\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4})$ in this specific case is chosen in such a way that the relative displacements of Ge and Te are equal in size but in opposite direction along [111].

The inverse transformation matrix–column pair $(\mathbf{Q}, \mathbf{q}) = (\mathbf{P}, \mathbf{p})^{-1} = (\mathbf{P}^{-1}, -\mathbf{P}^{-1}\mathbf{p})$ is necessary for the calculation of the atomic coordinates of the reference description X'_c . Given the matrix \mathbf{P} , its inverse \mathbf{P}^{-1} can be calculated either directly (*i.e.* applying the algebraic procedure for inversion of a matrix) or using the inverse matrices $\mathbf{Q}_1 = \mathbf{P}_1^{-1}$ and $\mathbf{Q}_2 = \mathbf{P}_2^{-1}$ listed in Table 1.5.1.1:

$$\begin{aligned} \mathbf{Q} = \mathbf{P}^{-1} &= (\mathbf{P}_1\mathbf{P}_2)^{-1} = \mathbf{P}_2^{-1}\mathbf{P}_1^{-1} \\ &= \begin{pmatrix} \frac{2}{3} & -\frac{1}{3} & -\frac{1}{3} \\ \frac{1}{3} & \frac{1}{3} & -\frac{2}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix} \begin{pmatrix} \bar{1} & 1 & 1 \\ 1 & \bar{1} & 1 \\ 1 & 1 & \bar{1} \end{pmatrix} = \begin{pmatrix} -\frac{4}{3} & \frac{2}{3} & \frac{2}{3} \\ -\frac{2}{3} & -\frac{2}{3} & \frac{4}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix}. \end{aligned} \quad (1.5.2.22)$$

(Note the change in the order of multiplication of the matrices \mathbf{P}_1^{-1} and \mathbf{P}_2^{-1} in \mathbf{Q} .) The corresponding origin shift \mathbf{q} is given by

$$\mathbf{q} = -\mathbf{P}^{-1}\mathbf{p} = -\begin{pmatrix} -\frac{4}{3} & \frac{2}{3} & \frac{2}{3} \\ -\frac{2}{3} & -\frac{2}{3} & \frac{4}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix} \begin{pmatrix} -\frac{1}{4} \\ -\frac{1}{4} \\ -\frac{1}{4} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \frac{1}{4} \end{pmatrix}. \quad (1.5.2.23)$$

The atomic positions of the reference description become

$$\begin{pmatrix} x'_c \\ y'_c \\ z'_c \end{pmatrix} = \begin{pmatrix} -\frac{4}{3} & \frac{2}{3} & \frac{2}{3} \\ -\frac{2}{3} & -\frac{2}{3} & \frac{4}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix} \begin{pmatrix} x_c \\ y_c \\ z_c \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \frac{1}{4} \end{pmatrix}.$$

The coordinates of the representative Ge atom occupying position $4a$ $0, 0, 0$ in $Fm\bar{3}m$ are transformed to $0, 0, \frac{1}{4}$, while those of Te are transformed from $4b$ $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ in $Fm\bar{3}m$ to $0, 0, \frac{3}{4}$. The comparison of these values with the experimentally determined atomic coordinates of Ge $0, 0, 0.2376$ and Te $0, 0, 0.7624$ reveals the corresponding atomic displacements associated with the displacive phase transition. The low-symmetry phase is a result of relative atomic displacements of the Ge and Te atoms along the polar (rhombohedral) [111] direction, giving rise to non-zero polarization along the same direction, *i.e.* the phase transition is a *paraelectric-to-ferroelectric* one.

1.5.3. Transformations between different space-group descriptions

BY G. CHAPUIS, H. WONDRAUSCHEK AND M. I. AROYO

1.5.3.1. Space groups with more than one description in this volume

In the description of the space-group symbols presented in Section 1.4.1, we have already seen that in the conventional,




Figure 1.5.3.1

Three possible cell choices for the monoclinic space group $P2_1/c$ (14) with unique axis b . Note the corresponding changes in the full Hermann–Mauguin symbols. The glide vector is indicated by an arrow.

unique axis b description of monoclinic space groups, the unique symmetry direction is chosen as \mathbf{b} ; it is normal to \mathbf{c} and \mathbf{a} , which form the angle β . However, it is often the case that this standard direction is not the most appropriate choice and that another choice would be more convenient. An example of this would be when following a phase transition from an orthorhombic parent phase to a monoclinic phase. Here, it would often be preferable to keep the same orientation of the axes even if the resulting monoclinic setting is not standard.

In some of the space groups, and especially in the monoclinic ones, the space-group tables of Chapter 2.3 provide a selection of possible alternative settings. For example, in space group $P2_1/c$, two possible orientations of the unit-cell axes are provided, namely with unique axis b and c . This is reflected in the corresponding full Hermann–Mauguin symbols by the explicit specification of the unique-axis position (dummy indices ‘1’ indicate ‘empty’ symmetry directions), and by the corresponding change in the direction of the glide plane: $P12_1/c1$ or $P112_1/a$ (*cf.* Section 1.4.1 for a detailed treatment of Hermann–Mauguin symbols of space groups).

It is not just the unique monoclinic axis that can be varied: the choice of the other axes can vary as well. There are cases where the selection of the conventional setting leads to an inconvenient monoclinic angle that deviates greatly from 90° . If another cell choice minimizes the deviation from 90° , it is preferred. Fig. 1.5.3.1 illustrates three cell choices for the monoclinic axis b setting of $P2_1/c$.

In centrosymmetric space groups the origin of the unit cell is located at an inversion centre (‘origin choice 2’). If, however, another point has higher site symmetry \mathcal{S} , a second diagram is displayed with the origin at a point with site symmetry \mathcal{S} (‘origin choice 1’). Fig. 1.5.3.2 illustrates the space group $Pban$ with two possible origins. The origin of the first choice is located on a point with site symmetry 222 , whereas the origin for the second choice is located on an inversion centre. Among the 230 space groups, this volume lists 24 centrosymmetric space groups with an additional alternative origin.

Finally, the seven rhombohedral space-group types (*i.e.* space groups with a rhombohedral lattice) also have alternative descriptions included in the space-group tables of this volume. The rhombohedral lattice is first presented with an R -centred