

1. GENERAL RELATIONSHIPS AND TECHNIQUES

three of them do not appear and the length of the others depends on the boundary plane, see Tables and Figs. 1.5.5.5 to 1.5.5.7.

The boundary conditions for the asymmetric unit are independent of the lattice parameters and the boundary plane is always represented by the simple equation $x, y, \frac{1}{2}; 0 < x, y < \frac{1}{2}$. By introducing flagpoles and wings, the description may become uni-arm.

1.5.5.4.2. Splitting of **k**-vector types

The Brillouin zone as well as the unit cell are always convex bodies; the same holds for the representation domain of CDML and for the choice of the asymmetric unit. It is thus sometimes unavoidable that the **k**-vector types are split and that the different parts belong to different arms and to different stars of **k** vectors. Sometimes this splitting of **k**-vector types may be avoided by an appropriate choice of the asymmetric unit; sometimes the introduction of flagpoles and wings is necessary to make the **k**-vector types uni-arm.

Examples

(1) In the reciprocal-space group $(\mathcal{G})^* = (Fm\bar{3}m)^*$, No. 225 of the arithmetic crystal class $m\bar{3}mI$ there are the lines of **k** vectors $\Lambda (\alpha, \alpha, \alpha)$ and $F (\frac{1}{2} - \alpha, -\frac{1}{2} + 3\alpha, \frac{1}{2} - \alpha)$ of CDML, p. 41. By Figure 1.5.5.1 one sees that the line Λ connects the points Γ and P , the line F connects the points P and H . One takes from the corresponding Table 1.5.5.1 the coefficients of $P = \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ and $H = 0, \frac{1}{2}, 0$. From these points or from the transformation listed at the top of Table 1.5.5.1 as 'Parameter relations' the coefficients of the line F are obtained as $F = x, \frac{1}{2} - x, x; 0 < x < \frac{1}{4}$.

The inspection of the symmetry diagram of $Fm\bar{3}m$, No. 225, in $IT A$ shows that a twofold rotation 2 (represented by the $4_2 \frac{1}{4}, y, \frac{1}{4}$ screw-rotation axis) leaves the point P invariant, whereas the point H is mapped onto the point $R \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$. More formally: the rotation is described by $x, \frac{1}{2} - x, x \rightarrow \frac{1}{2} - x, \frac{1}{2} - x, \frac{1}{2} - x$, where $0 < x < \frac{1}{4}$. The result is the line $F_1 = [RP]$. It is uni-arm to the line $\Lambda = x, x, x$ and the union $\Lambda \cup F_1$ forms the Wintgen position $32 f 3m$. An analogous result is obtained for the same lines in the arithmetic crystal class $m\bar{3}I$.

(2) In the following example the splitting of a Wintgen position happens if a representation domain of the Brillouin zone is chosen. The splitting can be avoided by the choice of the asymmetric unit. We consider the plane $x, y, 0$ in the arithmetic crystal class $4/mmmI$, see Fig. 1.5.5.4 and Table 1.5.5.4. In CDML this plane is split into the parts $C = [\Gamma S_2 R X]$ and $D = [M S G] \sim [M_2 S_2 R]$. By the choice of the asymmetric unit the independent region of the Wintgen position is uni-arm: $[\Gamma M_2 X] = 16 l m.. x, y, 0; 0 < x < y < \frac{1}{2}$.

(3) The splitting of a Wintgen position can be avoided if flagpoles and wings are admitted, i.e. if the minimal domain is described by a non-convex body. If one chooses in the first example of the arithmetic crystal classes $m\bar{3}mI$ and $m\bar{3}I$ the union $\Lambda \cup F_1$ for the line x, x, x , then $F_1 = [PR]$ forms a flagpole, whereas Λ forms an edge of the asymmetric unit, see Figs. 1.5.5.1 and 1.5.5.2.

The same holds for the Wintgen position $96 k ..m x, x, z$ of $m\bar{3}mI$. In the representation domain which is simultaneously the asymmetric unit, this Wintgen position is split into three parts B, C and J , which form three of the four walls of the (tetrahedral) minimal domain. By proper symmetry operations these three parts can be made uni-arm to the part C , such that their union $C \cup B_1 \cup J_1$ describes the independent part of that Wintgen position, see Fig. 1.5.5.1. The part C forms a wall of the asymmetric unit; the part $B_1 \cup J_1$ forms a wing, see Fig. 1.5.5.1.

1.5.5.4.3. **k**-vector types for non-holosymmetric space groups

The **k**-vector labels of CDML are primarily listed for the holosymmetric space groups. These lists are kept and supplemented for the non-holosymmetric space groups. In this way many superfluous **k**-vector labels are introduced.

Examples

(1) Arithmetic crystal class $m\bar{3}I$. In its reciprocal-space group $(Fm\bar{3})^*$, the introduction of the plane $AA = [\Gamma H_2 N]$ is unnecessary because the plane $A = [\Gamma N H]$ of Wintgen position $96 j m..$ of $(Fm\bar{3}m)^*$ can be extended to $A \cup AA = [\Gamma H_2 H]$ in the reciprocal-space group $(Fm\bar{3})^*$, cf. Fig. 1.5.5.2 and Table 1.5.5.2. In $(Fm\bar{3})^*$, both planes, A and AA , belong to Wintgen position $48 h m..$ The parameter description is extended from $x, y, 0; 0 < x < y < \frac{1}{2} - x (< \frac{1}{4})$ to $0 < y < \frac{1}{2} - x < \frac{1}{2}$.

(2) In the previous example, during the transition from the group $(Fm\bar{3}m)^*$ to the subgroup $(Fm\bar{3})^*$ the order of the little co-group of the special **k** vectors of $(Fm\bar{3}m)^*$ was not changed. In other cases, the little co-group may be reduced to a subgroup. Such **k** vectors may then be incorporated into a more general Wintgen position and described by an extension of the parameter range.

Arithmetic crystal class $m\bar{3}mI$, plane $[\Gamma H N] = x, y, 0$. In $(Fm\bar{3}m)^*$, see Fig. 1.5.5.1, all points (Γ, H, N) and lines (Δ, Σ, G) of the boundary of the asymmetric unit are special. In $(Fm\bar{3})^*$, see Fig. 1.5.5.2, the lines Δ and $[\Gamma H_2] \sim \Delta$ are special but Σ, G and $[N H_2] \sim G$ belong to the plane $(A \cup AA)$. The free parameter range on the line G is $0 < y < \frac{1}{4}$. Therefore, the parameter ranges of $(A \cup AA \cup G \cup \Sigma)$ in $x, y, 0$ can be taken as: $0 < x < \frac{1}{2} - y < \frac{1}{2}$ for $A \cup AA \cup \Sigma$ and $0 < y = \frac{1}{2} - x < \frac{1}{4}$ for G .

1.5.5.4.4. Ranges of independent parameters

In Section 1.5.4.3 a method for the determination of the parameter ranges was described. A few examples shall display the procedure.

(1) Arithmetic crystal class $m\bar{3}mI$, line $\Lambda \cup F_1$: In the reciprocal-space group $(Fm\bar{3}m)^*$ of the arithmetic crystal class $m\bar{3}mI$, the line x, x, x has stabilizer $\bar{3}m$ and little co-group $\bar{\mathcal{G}}^k = 3m$. Therefore, the divisor is $12:6 = 2$ and x is running from 0 to $\frac{1}{2}$.

The same result holds for the line $\Lambda \cup F_1$ in the reciprocal-space group $(Fm\bar{3})^*$ of the arithmetic crystal class $m\bar{3}I$: the stabilizer generated by $\bar{3}$ is of order 6, $|\bar{\mathcal{G}}^k| = |\{3\}| = 3$, the quotient is again $\frac{1}{2}$, the parameter range is the same as for $(Fm\bar{3}m)^*$.

(2) Arithmetic crystal class $m\bar{3}mI$, plane $B_1 \cup C \cup J_1$: In $(Fm\bar{3}m)^*$, the stabilizer of x, x, z is generated by $m.mm$ and the centring translation $t(\frac{1}{2}, \frac{1}{2}, 0)$ mod (integer translations). They generate a group of order 16; $\bar{\mathcal{G}}^k$ is $..m$ of order 2. The fraction of the plane is $\frac{2}{16} = \frac{1}{8}$ of the area $\sqrt{2}a^*2$ in the (centred) unit cell, as expressed by the parameter ranges $0 < x < \frac{1}{4}, 0 < z < \frac{1}{2}$. There are six arms of the star of x, x, z : $x, x, z; \bar{x}, x, z; x, y, x; x, y, \bar{x}; x, y, y; x, \bar{y}, y$. Three of them (x, x, z, \bar{x}, x, z and x, y, x) are represented in the boundaries of the representation domain: $C = [\Gamma N P], B = [H N P]$ and $J = [\Gamma H P]$, see Fig. 1.5.5.1. The areas of their parameter ranges are $\frac{1}{32}, \frac{1}{32}$ and $\frac{1}{16}$, respectively; the sum is $\frac{1}{8}$.

Arithmetic crystal class $m\bar{3}I$, the same result holds in the reciprocal-space group $(Fm\bar{3})^*$. The stabilizer generated by $2/m..$ and by the centring translation $t(\frac{1}{2}, \frac{1}{2}, 0)$ mod (integer translations) forms a group of order 8; the order of the little co-group $|\bar{\mathcal{G}}^k| = |\{1\}| = 1$. The quotient is again $\frac{1}{8}$, the parameter range is the same as for $(Fm\bar{3}m)^*$ but the plane belongs to the general position GP because the little co-group is trivial.

(3) Arithmetic crystal class $m\bar{3}mI$, reciprocal-space group $(Fm\bar{3}m)^*$, plane $x, y, 0$: the stabilizer of the plane A is generated by $4/mmm$ and $t(\frac{1}{2}, \frac{1}{2}, 0)$, order 32, $\bar{\mathcal{G}}^k$ (site-symmetry group) $m..$,