

2.4. ISOMORPHOUS REPLACEMENT AND ANOMALOUS SCATTERING

$$P(\alpha) = \prod P_s(\alpha) = N \exp \left(\sum_s K_s + \sum_s A_s \cos \alpha + \sum_s B_s \sin \alpha + \sum_s C_s \cos 2\alpha + \sum_s D_s \sin 2\alpha \right), \quad (2.4.4.36)$$

where K_s, A_s etc. are the constants appropriate for the s th source and N is the normalization constant.

2.4.4.8. Fourier representation of anomalous scatterers

It is often useful to have a Fourier representation of only the anomalous scatterers in a protein. The imaginary component of the electron-density distribution obviously provides such a representation. When the structure is known and $F_N(+)$ and $F_N(-)$ have been experimentally determined, Chacko & Srinivasan (1970) have shown that this representation is obtained in a Fourier synthesis with $i[F_N(+) + F_N^*(-)]/2$ as coefficients, where $F_N^*(-)$, whose magnitude is $F_N(-)$, is the complex conjugate of $F_N(+)$. They also indicated a method for calculating the phase angles of $F_N(+)$ and $F_N^*(-)$. It has been shown (Hendrickson & Sheriff, 1987) that the Bijvoet-difference Fourier synthesis proposed earlier by Kraut (1968) is an approximation of the true imaginary component of the electron density. The imaginary synthesis can be useful in identifying minor anomalous-scattering centres when the major centres are known and also in providing an independent check on the locations of anomalous scatterers and in distinguishing between anomalous scatterers with nearly equal atomic numbers (Sheriff & Hendrickson, 1987; Kitagawa *et al.*, 1987).

2.4.5. Anomalous scattering of neutrons and synchrotron radiation. The multiwavelength method

The multiwavelength anomalous-scattering method (Ramaseshan, 1982) relies on the variation of dispersion-correction terms as a function of the wavelength used. The success of the method therefore depends upon the size of the correction terms and the availability of incident beams of comparable intensities at different appropriate wavelengths. Thus, although this method was used as early as 1957 (Ramaseshan *et al.*, 1957) as an aid to structure solution employing characteristic X-rays, it is, as outlined below, ideally suited in structural work employing neutrons and synchrotron radiation. In principle, γ -radiation can also be used for phase determination (Raghavan, 1961; Moon, 1961) as the anomalous-scattering effects in γ -ray scattering could be very large; the wavelength is also easily tunable. However, the intensity obtainable for γ -rays is several orders lower than that obtainable from X-ray and neutron sources, and hence γ -ray anomalous scattering is of hardly any practical value in structural analysis.

2.4.5.1. Neutron anomalous scattering

Apart from the limitations introduced by experimental factors, such as the need for large crystals and the comparatively low flux of neutron beams, there are two fundamental reasons why neutrons are less suitable than X-rays for the *ab initio* determination of crystal structures. First, the neutron scattering lengths of different nuclei have comparable magnitudes whereas the atomic form factors for X-rays vary by two orders of magnitude. Therefore, Patterson techniques and the related heavy-atom method are much less suitable for use with neutron diffraction data than with X-ray data. Secondly, neutron scattering lengths could be positive or negative and hence, in general, the positivity criterion (Karle & Hauptman, 1950) or the squarability criterion (Sayre, 1952) does not hold good for nuclear density. Therefore,

the direct methods based on these criteria are not strictly applicable to structure analysis using neutron data, although it has been demonstrated that these methods could be successfully used in favourable situations in neutron crystallography (Sikka, 1969). The anomalous-scattering method is, however, in principle more powerful in the neutron case than in the X-ray case for *ab initio* structure determination.

Thermal neutrons are scattered anomalously at appropriate wavelengths by several nuclei. In a manner analogous to (2.4.3.1), the neutron scattering length of these nuclei can be written as

$$b_0 + b' + ib'' = b + ib''. \quad (2.4.5.1)$$

The correction terms b' and b'' are strongly wavelength-dependent. In favourable cases, b'/b_0 and b''/b_0 can be of the order of 10 whereas they are small fractions in X-ray anomalous scattering. In view of this pronounced anomalous effect in neutron scattering, Ramaseshan (1966) suggested that it could be used for structure solution. Subsequently, Singh & Ramaseshan (1968) proposed a two-wavelength method for unique structure analysis using neutron diffraction. The first part of the method is the determination of the positions of the anomalous scatterers from the estimated values of F_Q . The method employed for estimating F_Q is analogous to that using (2.4.4.9) except that data collected at two appropriate wavelengths are used instead of those from two isomorphous crystals. The second stage of the two-wavelength method involves phase evaluation. Referring to Fig. 2.4.3.2 and in a manner analogous to (2.4.3.5), we have

$$\sin \psi_1 = \frac{F_{N1}^2(+) - F_{N1}^2(-)}{4F_{N1}F_{Q1}'}, \quad (2.4.5.2)$$

where $\psi = \alpha_N - \alpha_Q$ and subscript 1 refers to data collected at wavelength λ_1 . Singh and Ramaseshan showed that $\cos \psi_1$ can also be determined when data are available at wavelength λ_1 and λ_2 . We may define

$$F_m^2 = [F_N^2(+) + F_N^2(-)]/2 \quad (2.4.5.3)$$

and we have from (2.4.3.3), (2.4.3.4) and (2.4.5.3)

$$F_N = (F_m^2 - F_Q^2)^{1/2}. \quad (2.4.5.4)$$

Then,

$$\cos \psi_1 = \frac{F_{m1}^2 - F_{m2}^2 - [(b_1^2 + b_1'^2) - (b_2^2 + b_2'^2)]x^2}{2(b_1 - b_2)F_{N1}x} + \frac{F_{Q1}}{F_{N1}}, \quad (2.4.5.5)$$

where x is the magnitude of the temperature-corrected geometrical part of F_Q . ψ_1 and hence α_{N1} can be calculated using (2.4.5.2) and (2.4.5.5). α_{N2} can also be obtained in a similar manner.

During the decade that followed Ramaseshan's suggestion, neutron anomalous scattering was used to solve half a dozen crystal structures, employing the multiple-wavelength methods as well as the methods developed for structure determination using X-ray anomalous scattering (Koetzle & Hamilton, 1975; Sikka & Rajagopal, 1975; Flook *et al.*, 1977). It has also been demonstrated that measurable Bijvoet differences could be obtained, in favourable situations, in neutron diffraction patterns from protein crystals (Schoenborn, 1975). However, despite the early promise held by neutron anomalous scattering, the method has

2. RECIPROCAL SPACE IN CRYSTAL-STRUCTURE DETERMINATION

not been as successful as might have been hoped. In addition to the need for large crystals, the main problem with using this method appears to be the time and expense involved in data collection (Koetzle & Hamilton, 1975).

2.4.5.2. Anomalous scattering of synchrotron radiation

The most significant development in recent years in relation to anomalous scattering of X-rays has been the advent of synchrotron radiation (Helliwell, 1984). The advantage of using synchrotron radiation for making anomalous-scattering measurements essentially arises out of the tunability of the wavelength. Unlike the characteristic radiation from conventional X-ray sources, synchrotron radiation has a smooth spectrum and the wavelength to be used can be finely selected. Accurate measurements have shown that values in the neighbourhood of 30 electrons could be obtained in favourable cases for f' and f'' (Templeton, Templeton, Phillips & Hodgson, 1980; Templeton, Templeton & Phizackerley, 1980; Templeton *et al.*, 1982). Schemes for the optimization of the wavelengths to be used have also been suggested (Narayan & Ramaseshan, 1981). Interestingly, the anomalous differences obtainable using synchrotron radiation are comparable in magnitude to the isomorphous differences normally encountered in protein crystallography. Thus, the use of anomalous scattering at several wavelengths would obviously eliminate the need for employing many heavy-atom derivatives. The application of anomalous scattering of synchrotron radiation for macromolecular structure analysis began to yield encouraging results in the 1980s (Helliwell, 1985). Intensity measurements from macromolecular X-ray diffraction patterns using synchrotron radiation at first relied primarily upon oscillation photography (Arndt & Wonacott, 1977). This method is not particularly suitable for accurately evaluating anomalous differences. Much higher levels of accuracy began to be achieved with the use of position-sensitive detectors (Arndt, 1986). Anomalous scattering, in combination with such detectors, has developed into a major tool in macromolecular crystallography (see *IT F*, 2001).

One of us (MV) acknowledges the support of the Department of Science & Technology, India.

References

- Adams, M. J. (1968). DPhil thesis, Oxford University, England.
- Agard, D. A. & Stroud, R. M. (1982). α -Bungarotoxin structure revealed by a rapid method for averaging electron density of non-crystallographically translationally related molecules. *Acta Cryst.* **A38**, 186–194.
- Arndt, U. W. (1986). X-ray position-sensitive detectors. *J. Appl. Cryst.* **19**, 145–163.
- Arndt, U. W. & Wonacott, A. J. (1977). *The Rotation Method in Crystallography*. Amsterdam: North-Holland.
- Ashida, T. (1976). Some remarks on the phase angle determination by the isomorphous replacement method. In *Crystallographic Computing Techniques*, edited by F. R. Ahmed, pp. 282–284. Copenhagen: Munksgaard.
- Bhat, T. N. & Blow, D. M. (1982). A density-modification method for improvement of poorly resolved protein electron-density maps. *Acta Cryst.* **A38**, 21–29.
- Bijvoet, J. M. (1949). Phase determination in direct Fourier-synthesis of crystal structures. *Proc. K. Ned. Akad. Wet. (B)*, **52**, 313–314.
- Bijvoet, J. M. (1954). Structure of optically active compounds in the solid state. *Nature (London)*, **173**, 888–891.
- Bijvoet, J. M., Peerdeman, A. F. & van Bommel, A. J. (1951). Determination of the absolute configuration of optically active compounds by means of X-rays. *Nature (London)*, **168**, 271–272.
- Blow, D. M. & Crick, F. H. C. (1959). The treatment of errors in the isomorphous replacement method. *Acta Cryst.* **12**, 794–802.
- Blow, D. M. & Matthews, B. W. (1973). Parameter refinement in the multiple isomorphous-replacement method. *Acta Cryst.* **A29**, 56–62.
- Blow, D. M. & Rossmann, M. G. (1961). *The single isomorphous replacement method*. *Acta Cryst.* **14**, 1195–1202.
- Blundell, T. L. & Johnson, L. N. (1976). *Protein Crystallography*. London: Academic Press.
- Bokhoven, C., Schoone, J. C. & Bijvoet, J. M. (1951). *The Fourier synthesis of the crystal structure of strychnine sulphate pentahydrate*. *Acta Cryst.* **4**, 275–280.
- Bradley, A. J. & Rodgers, J. W. (1934). *The crystal structure of the Heusler alloys*. *Proc. R. Soc. London Ser. A*, **144**, 340–359.
- Cannillo, E., Oberti, R. & Ungaretti, L. (1983). Phase extension and refinement by density modification in protein crystallography. *Acta Cryst.* **A39**, 68–74.
- Chacko, K. K. & Srinivasan, R. (1970). On the Fourier refinement of anomalous dispersion corrections in X-ray diffraction data. *Z. Kristallogr.* **131**, 88–94.
- Collins, D. M. (1975). Efficiency in Fourier phase refinement for protein crystal structures. *Acta Cryst.* **A31**, 388–389.
- Cork, J. M. (1927). *The crystal structure of some of the alums*. *Philos. Mag.* **4**, 688–698.
- Coster, D., Knol, K. S. & Prins, J. A. (1930). Unterschiede in der Intensität der Röntgenstrahlenreflexion an den beiden 111-Flächen der Zinkblende. *Z. Phys.* **63**, 345–369.
- Cromer, D. T. (1965). Anomalous dispersion corrections computed from self-consistent field relativistic Dirac-Slater wave functions. *Acta Cryst.* **18**, 17–23.
- Cruickshank, D. W. J. & McDonald, W. S. (1967). Parameter errors in polar space groups caused by neglect of anomalous scattering. *Acta Cryst.* **23**, 9–11.
- Cullis, A. F., Muirhead, H., Perutz, M. F., Rossmann, M. G. & North, A. C. T. (1961a). *The structure of haemoglobin. VIII. A three-dimensional Fourier synthesis at 5.5 Å resolution: determination of the phase angles*. *Proc. R. Soc. London Ser. A*, **265**, 15–38.
- Cullis, A. F., Muirhead, H., Perutz, M. F., Rossmann, M. G. & North, A. C. T. (1961b). *The structure of haemoglobin. IX. A three-dimensional Fourier synthesis at 5.5 Å resolution: description of the structure*. *Proc. R. Soc. London Ser. A*, **265**, 161–187.
- Dale, D., Hodgkin, D. C. & Venkatesan, K. (1963). *The determination of the crystal structure of factor V Ia*. In *Crystallography and Crystal Perfection*, edited by G. N. Ramachandran, pp. 237–242. New York, London: Academic Press.
- Dickerson, R. E., Kendrew, J. C. & Strandberg, B. E. (1961). *The crystal structure of myoglobin: phase determination to a resolution of 2 Å by the method of isomorphous replacement*. *Acta Cryst.* **14**, 1188–1195.
- Dickerson, R. E., Weinzierl, J. E. & Palmer, R. A. (1968). A least-squares refinement method for isomorphous replacement. *Acta Cryst.* **B24**, 997–1003.
- Dodson, E., Evans, P. & French, S. (1975). *The use of anomalous scattering in refining heavy atom parameters in proteins*. In *Anomalous Scattering*, edited by S. Ramaseshan & S. C. Abrahams, pp. 423–436. Copenhagen: Munksgaard.
- Dodson, E. & Vijayan, M. (1971). *The determination and refinement of heavy-atom parameters in protein heavy-atom derivatives. Some model calculations using acentric reflexions*. *Acta Cryst.* **B27**, 2402–2411.
- Einstein, J. E. (1977). *An improved method for combining isomorphous replacement and anomalous scattering diffraction data for macromolecular crystals*. *Acta Cryst.* **A33**, 75–85.
- Flook, R. J., Freeman, H. C. & Scudder, M. L. (1977). *An X-ray and neutron diffraction study of aqua(L-glutamato) cadmium(II) hydrate*. *Acta Cryst.* **B33**, 801–809.
- Gilli, G. & Cruickshank, D. W. J. (1973). Effect of neglect of dispersion in centrosymmetric structures: results for OsO_4 . *Acta Cryst.* **B29**, 1983–1985.
- Green, D. W., Ingram, V. M. & Perutz, M. F. (1954). *The structure of haemoglobin. IV. Sign determination by the isomorphous replacement method*. *Proc. R. Soc. London Ser. A*, **225**, 287–307.
- Green, E. A. (1979). A new statistical model for describing errors in isomorphous replacement data: the case of one derivative. *Acta Cryst.* **A35**, 351–359.
- Harker, D. (1956). *The determination of the phases of the structure factors of non-centrosymmetric crystals by the method of double isomorphous replacement*. *Acta Cryst.* **9**, 1–9.
- Helliwell, J. R. (1984). *Synchrotron X-radiation protein crystallography: instrumentation, methods and applications*. *Rep. Prog. Phys.* **47**, 1403–1497.