

4. DIFFUSE SCATTERING AND RELATED TOPICS




Fig. 4.6.3.15. Atomic surface of the Penrose tiling in the 5D hypercubic description. The projection of the 5D hypercubic unit cell upon \mathbf{V}^\perp gives a rhomb-icosahedron (above). The Penrose tiling is generated by four equidistant pentagons (shaded) inscribed in the rhomb-icosahedron. Below is a perpendicular-space projection of the same pentagons, which are located on the $[1111]_D$ diagonal of the 4D hyper-rhombohedral unit cell in the 4D description.

$$W = \begin{pmatrix} a_1^* \cos(2\pi/5) & a_2^* \cos(4\pi/5) & a_3^* \cos(6\pi/5) & a_4^* \cos(8\pi/5) \\ a_1^* \sin(2\pi/5) & a_2^* \sin(4\pi/5) & a_3^* \sin(6\pi/5) & a_4^* \sin(8\pi/5) \\ a_1^* \cos(4\pi/5) & a_2^* \cos(8\pi/5) & a_3^* \cos(2\pi/5) & a_4^* \cos(6\pi/5) \\ a_1^* \sin(4\pi/5) & a_2^* \sin(8\pi/5) & a_3^* \sin(2\pi/5) & a_4^* \sin(6\pi/5) \end{pmatrix}.$$

The 2D Penrose tiling can also be embedded canonically in the 5D space. Canonically means that the 5D lattice is hypercubic and that the projection of one unit cell upon the 3D perpendicular space \mathbf{V}^\perp , giving a rhomb-icosahedron, defines the atomic surface. However, the parallel-space image \mathbf{a}_i^* , $i = 1, \dots, 4$, with $\mathbf{a}_0^* = -(\mathbf{a}_1^* + \mathbf{a}_2^* + \mathbf{a}_3^* + \mathbf{a}_4^*)$, of the 5D basis \mathbf{d}_i^* , $i = 1, \dots, 4$ is not linearly independent. Consequently, the atomic surface consists of only a subset of the points contained in the rhomb-icosahedron: five equidistant pentagons (one with diameter zero) resulting as sections of the rhomb-icosahedron with five equidistant parallel planes (Fig. 4.6.3.15). The linear dependence of the 5D basis allows the embedding in the 4D space. The resulting hyper-rhombohedral hyperlattice is spanned by the basis \mathbf{d}_i , $i = 1, \dots, 4$, discussed above. The atomic surfaces occupy the positions $p/5(1111)$, $p = 1, \dots, 4$, on the body diagonal of the 4D unit cell. Neighbouring pentagons are in an *anti* position to each other (Fig. 4.6.3.16). Thus the 4D unit cell is decorated centrosymmetrically. The edge length a_r of a Penrose rhomb is related to the length of physical-space basis vectors a_i^* by $a_r = \tau S$, with the smallest distance $S = (2\tau/5a_i^*)$, $i = 1, \dots, 4$. The *point density* (number of vertices per unit area) of a Penrose tiling with Penrose rhombs of edge length a_r can be calculated from the ratio of the relative number of unit tiles in the tiling to their area:

$$\rho = \frac{1 + \tau}{a_r^2 [\sin(\pi/5) + \tau \sin(2\pi/5)]} = (5/2)a_i^{*2} (2 - \tau)^2 \tan(2\pi/5).$$

This is equivalent to the calculation from the 4D description,

$$\begin{aligned} \rho &= \frac{\sum_{i=1}^4 \Omega_{AS}^i}{\Omega_{UC}} = \frac{\sum_{i=1}^4 (5/2)\lambda^2 \sin(2\pi/5)}{4/[5(5)^{1/2}|a_i^*|^4]} \\ &= (5/2)a_i^{*2} (2 - \tau)^2 \tan(2\pi/5), \end{aligned}$$

where Ω_{AS} and Ω_{UC} are the area of the atomic surface and the volume of the 4D unit cell, respectively. The pentagon radii are




Fig. 4.6.3.16. Projection of the 4D hyper-rhombohedral unit cell of the Penrose tiling in the 4D description upon the perpendicular space. In the upper drawing all edges between the 16 corners are shown. In the lower drawing the corners are indexed and the four pentagonal atomic surfaces of the Penrose tiling are shaded.

$\lambda_{1,4} = 2(2 - \tau)/5a^*$ and $\lambda_{2,3} = 2(\tau - 1)/5a^*$ for the atomic surfaces in $(p/5)(1111)$ with $p = 1, 4$ and $p = 2, 3$. A detailed discussion of the properties of Penrose tiling is given in the papers of Penrose (1974, 1979), Jaric (1986) and Pavlovitch & Kleman (1987).

4.6.3.3.2.1. Indexing

The indexing of the submodule M_1^* of the diffraction pattern of a decagonal phase is not unique. Since M_1^* corresponds to a \mathbb{Z} module of rank 4 with decagonal point symmetry, it is invariant under scaling by τ^n , $n \in \mathbb{Z}$: $S^n M^* = \tau^n M^*$. Nevertheless, an optimum basis (low indices are assigned to strong reflections) can be derived: not the metrics, as for regular periodic crystals, but the intensity distribution characterizes the best choice of indexing.

A correct set of reciprocal-basis vectors can be identified experimentally in the following way:

(1) Find directions of systematic absences or pseudo-absences determining the possible orientations of the reciprocal-basis vectors (see Rabson *et al.*, 1991).

(2) Find pairs of strong reflections whose physical-space diffraction vectors are related to each other by the factor τ .

(3) Index these reflections by assigning an appropriate value to a^* . This value should be derived from the shortest interatomic distance S and the edge length of the unit tiles expected in the structure.

(4) The reciprocal basis is correct if all observable Bragg reflections can be indexed with integer numbers.

4.6.3.3.2.2. Diffraction symmetry

The diffraction symmetry of decagonal phases can be described by the Laue groups $10/mmm$ or $10/m$. The set of all vectors \mathbf{H} forms a Fourier module $M^* = \{\mathbf{H}^\parallel = \sum_{i=1}^5 h_i \mathbf{a}_i^* | h_i \in \mathbb{Z}\}$ of rank 5 in physical space which can be decomposed into two submodules $M^* = M_1^* \oplus M_2^*$. $M_1^* = \{h_1 \mathbf{a}_1^* + h_2 \mathbf{a}_2^* + h_3 \mathbf{a}_3^* + h_4 \mathbf{a}_4^*\}$ corresponds to a \mathbb{Z} module of rank 4 in a 2D subspace, $M_2^* = \{h_5 \mathbf{a}_5^*\}$ corresponds to a \mathbb{Z} module of rank 1 in a 1D subspace. Consequently, the first submodule can be considered as a projection from a 4D reciprocal lattice, $M_1^* = \pi^\parallel(\Sigma^*)$, while the second submodule is of the form of a regular 1D reciprocal lattice, $M_2^* = \Lambda^*$. The diffraction pattern of the Penrose tiling decorated with equal point scatterers on its vertices is shown in Fig. 4.6.3.17. All Bragg reflections within