

1. TENSORIAL ASPECTS OF PHYSICAL PROPERTIES

1.1.3.3.2. *Multiplication by a scalar*

This is a particular case of the tensor product.

1.1.3.3.3. *Contracted product, contraction*

Here we are concerned with an operation that only exists in the case of tensors and that is very important because of its applications in physics. In practice, it is almost always the case that tensors enter into physics through the intermediary of a contracted product.

(i) *Contraction*. Let us consider a tensor of rank 2 that is once covariant and once contravariant. Let us write its transformation in a change of coordinate system:

$$t_i^j = A_p^j B_i^q t_q^p.$$

Now consider the quantity t_i^i derived by applying the Einstein convention ($t_i^i = t_1^1 + t_2^2 + t_3^3$). It follows that

$$t_i^i = A_p^i B_i^q t_q^p = \delta_p^q t_q^p \\ t_i^i = t_p^p.$$

This is an invariant quantity and so is a *scalar*. This operation can be carried out on any tensor of rank higher than or equal to two, provided that it is expressed in a form such that its components are (at least) once covariant and once contravariant.

The *contraction* consists therefore of equalizing a covariant index and a contravariant index, and then in summing over this index. Let us take, for example, the tensor t_i^{jk} . Its contracted form is t_i^{ik} , which, with a change of basis, becomes

$$t_i^{ik} = A_p^k t_q^{ip}.$$

The components t_i^{ik} are those of a vector, resulting from the contraction of the tensor t_i^{jk} . The rank of the tensor has changed from 3 to 1. In a general manner, the contraction reduces the rank of the tensor from n to $n - 2$.

Example. Let us take again the operator of symmetry O . The trace of the associated matrix is equal to

$$O_1^1 + O_2^2 + O_3^3 = O_i^i.$$

It is the resultant of the contraction of the tensor O . It is a tensor of rank 0, which is a scalar and is invariant under a change of basis.

(ii) *Contracted product*. Consider the product of two tensors of which one is contravariant at least once and the other covariant at least once:

$$p_i^{jk} = t_i^j z^k.$$

If we contract the indices i and k , it follows that

$$p_i^j = t_i^j z^i.$$

The contracted product is then a tensor of rank 1 and not 3. It is an operation that is very frequent in practice.

(iii) *Scalar product*. Next consider the tensor product of two vectors:

$$t_i^j = x_i y^j.$$

After contraction, we get the scalar product:

$$t_i^i = x_i y^i.$$

1.1.3.4. *Tensor nature of physical quantities*

Let us first consider the dielectric constant. In the introduction, we remarked that for an isotropic medium

$$\mathbf{D} = \epsilon \mathbf{E}.$$

If the medium is anisotropic, we have, for one of the components,

$$D^1 = \epsilon_1^1 E^1 + \epsilon_2^1 E^2 + \epsilon_3^1 E^3.$$

This relation and the equivalent ones for the other components can also be written

$$D^i = \epsilon_j^i E^j \tag{1.1.3.3}$$

using the Einstein convention.

The scalar product of \mathbf{D} by an arbitrary vector \mathbf{x} is

$$D^i x_i = \epsilon_j^i E^j x_i.$$

The right-hand member of this relation is a bilinear form that is invariant under a change of basis. The set of nine quantities ϵ_j^i constitutes therefore the set of components of a tensor of rank 2. Expression (1.1.3.3) is the contracted product of ϵ_j^i by E^j .

A similar demonstration may be used to show the tensor nature of the various physical properties described in Section 1.1.1, whatever the rank of the tensor. Let us for instance consider the piezoelectric effect (see Section 1.1.4.4.3). The components of the electric polarization, P^i , which appear in a medium submitted to a stress represented by the second-rank tensor T_{jk} are

$$P^i = d^{ijk} T_{jk},$$

where the tensor nature of T_{jk} will be shown in Section 1.3.2. If we take the contracted product of both sides of this equation by any vector of covariant components x_i , we obtain a linear form on the left-hand side, and a trilinear form on the right-hand side, which shows that the coefficients d^{ijk} are the components of a third-rank tensor. Let us now consider the piezo-optic (or photoelastic) effect (see Sections 1.1.4.10.5 and 1.6.7). The components of the variation $\Delta\eta^{ij}$ of the dielectric impermeability due to an applied stress are

$$\Delta\eta^{ij} = \pi^{ijkl} T_{jl}.$$

In a similar fashion, consider the contracted product of both sides of this relation by two vectors of covariant components x_i and y_j , respectively. We obtain a bilinear form on the left-hand side, and a quadrilinear form on the right-hand side, showing that the coefficients π^{ijkl} are the components of a fourth-rank tensor.

1.1.3.5. *Representation surface of a tensor*

1.1.3.5.1. *Definition*

Let us consider a tensor $t_{ijkl\dots}$ represented in an orthonormal frame where variance is not important. The value of component $t'_{1111\dots}$ in an arbitrary direction is given by

$$t'_{1111\dots} = t_{ijkl\dots} B_1^i B_1^j B_1^k B_1^l \dots,$$

where the B_1^i, B_1^j, \dots are the direction cosines of that direction with respect to the axes of the orthonormal frame.

The *representation surface* of the tensor is the polar plot of $t'_{1111\dots}$.

1.1.3.5.2. *Representation surfaces of second-rank tensors*

The representation surfaces of second-rank tensors are quadrics. The directions of their principal axes are obtained as follows. Let t_{ij} be a second-rank tensor and let $\mathbf{OM} = \mathbf{r}$ be a vector with coordinates x_i . The doubly contracted product, $t_{ij} x^i x^j$, is a scalar. The locus of points M such that