

1. TENSORIAL ASPECTS OF PHYSICAL PROPERTIES

1.1.4.5.3.2. *Tensors of higher rank*

If the rank of the tensor is higher than 2, the tensor may be antisymmetric with respect to the indices of one or several couples of indices.

(i) *Tensors of rank 3 antisymmetric with respect to every couple of indices.* A trilinear form  $T(\mathbf{x}, \mathbf{y}, \mathbf{z}) = t_{ijk}x^i y^j z^k$  is said to be antisymmetric if it satisfies the relations

$$\left. \begin{aligned} T(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= -T(\mathbf{y}, \mathbf{x}, \mathbf{z}) \\ &= -T(\mathbf{x}, \mathbf{z}, \mathbf{y}) \\ &= -T(\mathbf{z}, \mathbf{y}, \mathbf{x}). \end{aligned} \right\}$$

Tensor  $t_{ijk}$  has 27 components. It is found that all of them are equal to zero, except

$$t_{123} = t_{231} = t_{312} = -t_{213} = -t_{132} = -t_{321}.$$

The three-times contracted product with the permutations tensor (Section 1.1.3.7.2),  $(1/6)\varepsilon_{ijk}t_{ijk}$ , is a *pseudoscalar* or *axial scalar*. It is not a usual scalar: the sign of this product changes when one changes the hand of the reference axes, change of basis represented by the matrix

$$\begin{pmatrix} \bar{1} & 0 & 0 \\ 0 & \bar{1} & 0 \\ 0 & 0 & \bar{1} \end{pmatrix}.$$

Form  $T(\mathbf{x}, \mathbf{y}, \mathbf{z})$  can also be written

$$T(\mathbf{x}, \mathbf{y}, \mathbf{z}) = P t_{123},$$

where

$$P = \varepsilon_{ijk}x^i y^j z^k = \begin{vmatrix} x^1 & x^2 & x^3 \\ y^1 & y^2 & y^3 \\ z^1 & z^2 & z^3 \end{vmatrix}$$

is the triple scalar product of the three vectors  $\mathbf{x}, \mathbf{y}, \mathbf{z}$ :

$$P = (\mathbf{x}, \mathbf{y}, \mathbf{z}) = (\mathbf{x} \wedge \mathbf{y} \cdot \mathbf{z}).$$

It is also a pseudoscalar. The permutation tensor is not a real tensor of rank 3: if the hand of the axes is changed, the sign of  $P$  also changes;  $P$  is therefore not a trilinear form.

Another example of a pseudoscalar is given by the rotatory power of an optically active medium, which is expressed through the relation (see Section 1.6.5.4)

$$\theta = \rho d,$$

where  $\theta$  is the rotation angle of the light wave,  $d$  the distance traversed in the material and  $\rho$  is a pseudoscalar: if one takes the mirror image of this medium, the sign of the rotation of the light wave also changes.

(ii) *Tensor of rank 3 antisymmetric with respect to one couple of indices.* Let us consider a trilinear form such that

$$T(\mathbf{x}, \mathbf{y}, \mathbf{z}) = -T(\mathbf{y}, \mathbf{x}, \mathbf{z}).$$

Its components satisfy the relation

$$t^{iil} = 0; \quad t^{ijl} = -t^{jil}.$$

The twice contracted product

$$t_k^l = \frac{1}{2} \varepsilon_{ijk} t^{ijl}$$

is an *axial* tensor of rank 2 whose components are the independent components of the antisymmetric tensor of rank 3,  $t^{ijl}$ .

*Examples*

(1) *Hall constant.* The Hall effect is observed in semiconductors. If one takes a semiconductor crystal and applies a

magnetic induction  $\mathbf{B}$  and at the same time imposes a current density  $\mathbf{j}$  at right angles to it, one observes an electric field  $\mathbf{E}$  at right angles to the other two fields (see Section 1.8.3.4). The expression for the field can be written

$$E_i = R_H \varepsilon_{ikl} j_k B_l,$$

where  $R_H \varepsilon_{ikl}$  is the Hall constant, which is a tensor of rank 3. However, because the direction of the current density is imposed by the physical law (the set of vectors  $\mathbf{B}, \mathbf{j}, \mathbf{E}$  constitutes a right-handed frame), one has

$$R_H \varepsilon_{ikl} = -R_H \varepsilon_{kil},$$

which shows that  $R_H \varepsilon_{ikl}$  is an antisymmetric (axial) tensor of rank 3. As can be seen from its physical properties, only the components such that  $i \neq k \neq l$  are different from zero. These are

$$R_H \varepsilon_{123} = -R_H \varepsilon_{213}; \quad R_H \varepsilon_{132} = -R_H \varepsilon_{312}; \quad R_H \varepsilon_{312}; \quad R_H \varepsilon_{321}.$$

(2) *Optical rotation.* The gyration tensor used to describe the property of optical rotation presented by gyrotropic materials (see Section 1.6.5.4) is an axial tensor of rank 2, which is actually an antisymmetric tensor of rank 3.

(3) *Acoustic activity.* The acoustic gyrotropic tensor describes the rotation of the polarization plane of a transverse acoustic wave propagating along the acoustic axis (see for instance Kumaraswamy & Krishnamurthy, 1980). The elastic constants may be expanded as

$$c_{ijkl}(\omega, \mathbf{k}) = c_{ijkl}(\omega) + id_{ijklm}(\omega)k_m + \dots,$$

where  $d_{ijklm}$  is a fifth-rank tensor. Time-reversal invariance requires that  $d_{ijklm} = -d_{jiklm}$ , which shows that it is an antisymmetric (axial) tensor.

1.1.4.5.3.3. *Properties of axial tensors*

The two preceding sections have shown examples of axial tensors of ranks 0 (pseudoscalar), 1 (pseudovector) and 2. They have in common that all their components change sign when the sign of the basis is changed, and this can be taken as the definition of an axial tensor. Their components are the components of an antisymmetric tensor of higher rank. It is important to bear in mind that in order to obtain their behaviour in a change of basis, one should first determine the behaviour of the components of this antisymmetric tensor.

1.1.4.6. *Symmetry of tensors imposed by the crystalline medium*

Many papers have been devoted to the derivation of the invariant components of physical property tensors under the influence of the symmetry elements of the crystallographic point groups: see, for instance, Fumi (1951, 1952a,b,c, 1987), Fumi & Ripamonti (1980a,b), Nowick (1995), Nye (1957, 1985), Sands (1995), Sirotnin & Shaskol'skaya (1982), and Wooster (1973). There are three main methods for this derivation: the matrix method (described in Section 1.1.4.6.1), the direct inspection method (described in Section 1.1.4.6.3) and the group-theoretical method (described in Section 1.2.4 and used in the accompanying software, see Section 1.2.7.4).

1.1.4.6.1. *Matrix method – application of Neumann's principle*

An operation of symmetry turns back the crystalline edifice on itself; it allows the physical properties of the crystal and the tensors representing them to be invariant. An operation of symmetry is equivalent to a change of coordinate system. In a change of system, a tensor becomes

$$t_{\gamma\delta}^{\alpha\beta} = t_{kl}^{ij} A_i^\alpha A_j^\beta B_\gamma^k B_\delta^l.$$

If  $A$  represents a symmetry operation, it is a unitary matrix:

## 1.1. INTRODUCTION TO THE PROPERTIES OF TENSORS

$$A = B^T = B^{-1}.$$

Since the tensor is invariant under the action of the symmetry operator  $A$ , one has, according to Neumann's principle,

$$t_{\gamma\delta}^{\alpha\beta} = t_{\gamma\delta}^{\alpha\beta}$$

and, therefore,

$$t_{\gamma\delta}^{\alpha\beta} = t_{kl}^{ij} A_i^\alpha A_j^\beta B_\gamma^k B_\delta^l. \quad (1.1.4.1)$$

There are therefore a certain number of linear relations between the components of the tensor and the number of independent components is reduced. If there are  $p$  components and  $q$  relations between the components, there are  $p - q$  independent components. This number is *independent* of the system of axes. When applied to each of the 32 point groups, this reduction enables one to find the form of the tensor in each case. It depends on the rank of the tensor. In the present chapter, the reduction will be derived for tensors up to the fourth rank and for all crystallographic groups as well as for the isotropic groups. An orthonormal frame will be assumed in all cases, so that co- and contravariance will not be apparent and the positions of indices as subscripts or superscripts will not be meaningful. The  $Ox_3$  axis will be chosen parallel to the threefold, fourfold or sixfold axis in the trigonal, tetragonal and hexagonal systems. The accompanying software to the present volume enables the reduction for tensors of any rank to be derived.

### 1.1.4.6.2. The operator $A$ is in diagonal form

#### 1.1.4.6.2.1. Introduction

If one takes as the system of axes the eigenvectors of the operator  $A$ , the matrix is written in the form

$$\begin{pmatrix} \exp i\theta & 0 & 0 \\ 0 & \exp -i\theta & 0 \\ 0 & 0 & \pm 1 \end{pmatrix},$$

where  $\theta$  is the rotation angle,  $Ox_3$  is taken parallel to the rotation axis and coefficient  $A_3$  is equal to +1 or -1 depending on whether the rotation axis is direct or inverse (proper or improper operator).

The equations (1.1.4.1) can then be simplified and reduce to

$$t_{kl}^{ij} = t_{kl}^{ij} A_i^\alpha A_j^\beta B_\gamma^k B_\delta^l \quad (1.1.4.2)$$

(without any summation).

If the product  $A_i^\alpha A_j^\beta B_\gamma^k B_\delta^l$  (without summation) is equal to unity, equation (1.1.4.2) is trivial and there is significance in the component  $t_{kl}$ . On the contrary, if it is different from 1, the only solution for (1.1.4.2) is that  $t_{kl}^{ij} = 0$ . One then finds immediately that certain components of the tensor are zero and that others are unchanged.

#### 1.1.4.6.2.2. Case of a centre of symmetry

All the diagonal components are in this case equal to -1. One thus has:

(i) *Tensors of even rank*,  $t^{ij\dots} = (-1)^{2p} t^{ij\dots}$ . The components are not affected by the presence of the centre of symmetry. The reduction of tensors of even rank is therefore the same in a centred group and in its noncentred subgroups, that is in any of the 11 *Laue classes*:

$\bar{1}$	1
$2/m$	2, $m$
$mmm$	222, $2mm$
$\bar{3}$	3
$\bar{3}m$	32, $3m$
$4/m$	4, $4$
$4/m\bar{m}$	422, $4mm$
$6/m$	6, $6$
$6/m\bar{m}$	622, $6mm$
$m\bar{3}$	23
$m\bar{3}m$	432, $\bar{4}32$ .

If a tensor is invariant with respect to two elements of symmetry, it is invariant with respect to their product. It is then sufficient to make the reduction for the generating elements of the group and (since this concerns a tensor of even rank) for the 11 *Laue classes*.

(ii) *Tensors of odd rank*,  $t^{ij\dots} = (-1)^{2p+1} t^{ij\dots}$ . All the components are equal to zero. The physical properties represented by tensors of rank 3, such as piezoelectricity, piezomagnetism, nonlinear optics, for instance, will therefore not be present in a centrosymmetric crystal.

#### 1.1.4.6.2.3. General case

By replacing the matrix coefficients  $A_i^\alpha$  by their expression, (1.1.4.2) becomes, for a proper rotation,

$$t^{jk\dots} = t^{jk\dots} \exp(ir\theta) \exp(-is\theta)(1)^t = t^{jk\dots} \exp i(r-s)\theta,$$

where  $r$  is the number of indices equal to 1,  $s$  is the number of indices equal to 2,  $t$  is the number of indices equal to 3 and  $r + s + t = p$  is the rank of the tensor. The component  $t^{jk\dots}$  is not affected by the symmetry operation if

$$(r-s)\theta = 2K\pi,$$

where  $K$  is an integer, and is equal to zero if

$$(r-s)\theta \neq 2K\pi.$$

The angle of rotation  $\theta$  can be put into the form  $2\pi/q$ , where  $q$  is the order of the axis. The condition for the component not to be zero is then

$$r-s = Kq.$$

The condition is fulfilled differently depending on the rank of the tensor,  $p$ , and the order of the axis,  $q$ . Indeed, we have  $r-s \leq p$  and

$p = 2, r-s \leq 2$ : the result of the reduction will be the same for any  $q \geq 3$ ;

$p = 3, r-s \leq 3$ : the result of the reduction will be the same for any  $q \geq 4$ ;

$p = 4, r-s \leq 4$ : the result of the reduction will be the same for any  $q \geq 5$ .

It follows that:

(i) for tensors of rank 2, the reduction will be the same for trigonal (threefold axis), tetragonal (fourfold axis) and hexagonal (sixfold axis) groups;

(ii) for tensors of rank 3, the reduction will be the same for tetragonal and hexagonal groups;

(iii) for tensors of rank 4, the reduction will be different for trigonal, tetragonal and hexagonal groups.

The inconvenience of the diagonalization method is that the vectors and eigenvalues are, in general, complex, so in practice one uses another method. For instance, we may note that equation (1.1.4.1) can be written in the case of  $p = 2$  by associating with the tensor a  $3 \times 3$  matrix  $T$ :

$$T = BTB^T,$$