


1. TENSORIAL ASPECTS OF PHYSICAL PROPERTIES

1.1.4.8.7.2. Groups 432 and $\infty A_{\infty}/M$


One combines the reductions corresponding to groups 422 and 23 :



There is 1 independent component.

1.1.4.8.7.3. Group $\bar{4}3m$

One combines the reductions corresponding to groups $\bar{4}2m$ and 23 :



There is 1 independent component.

1.1.4.8.7.4. Groups $m\bar{3}$, $m\bar{3}m$ and $\infty(A_{\infty}/M)C$

All the components are equal to zero.


1.1.4.9. Reduction of the components of a tensor of rank 4

1.1.4.9.1. Triclinic system (groups $\bar{1}$, 1)

There is no reduction; all the components are independent. Their number is equal to 81. They are usually represented as a 9×9 matrix, where components t_{ijkl} are replaced by $ijkl$, for brevity:

ij	kl	11	22	33	23	31	12	32	13	21
11	1111	1122	1133	1123	1131	1112	1132	1113	1121	1131
22	2211	2222	2233	2223	2231	2212	2232	2213	2221	2231
33	3311	3322	3333	3323	3331	3312	3332	3313	3321	3331
23	2311	2322	2333	2323	2331	2312	2332	2313	2321	2331
31	3111	3122	3133	3123	3131	3112	3132	3113	3121	3131
12	1211	1222	1233	1223	1231	1212	1232	1213	1221	1231
32	3211	3222	3233	3223	3231	3212	3232	3213	3221	3231
13	1311	1322	1333	1323	1331	1312	1332	1313	1321	1331
21	2111	2122	2133	2123	2131	2112	2132	2113	2121	2131


This matrix can be represented symbolically by



where the 9×9 matrix has been subdivided for clarity in to nine 3×3 submatrices.


1.1.4.9.2. Monoclinic system (groups $2/m$, 2 , m)

The reduction is obtained by the method of direct inspection. For a twofold axis parallel to Ox_2 , one finds



There are 41 independent components.

1.1.4.9.3. Orthorhombic system (groups mmm , $2mm$, 222)



There are 21 independent components.

1.1.4.9.4. Trigonal system

1.1.4.9.4.1. Groups 3 and $\bar{3}$

The reduction is first applied in the system of axes tied to the eigenvectors of the operator representing a threefold axis. The system of axes is then changed to a system of orthonormal axes with Ox_3 parallel to the threefold axis:

ij	kl	11	22	33	23	31	12	32	13	21
11	1111	1122	1133	1123	-2231	1112	1132	-2213	1121	1131
22	1122	1111	1133	-1123	2231	-1121	-1132	2213	-1121	-1131
33	3311	3311	3333			3312				-3312
23	2311	-2311		2323	2331	1322	2332	2313	1322	2331
31	-3122	3122		3123	3131	3211	3132	3113	3211	3131
12	1211	-2111	1233	2213	1132	1212	2231	1123	1221	2231
32	3211	-3211		3113	-3132	3122	3131	-3123	3122	3131
13	-1322	1322		-2313	2332	2311	-2331	2323	2311	2331
21	2111	-1211	-1233	2213	1132	1221	2231	1123	1221	2231

with

$$\left. \begin{aligned} t_{1111} - t_{1122} &= t_{1212} + t_{1221} \\ t_{1112} + t_{1121} &= -(t_{1211} + t_{2111}). \end{aligned} \right\}$$

There are 27 independent components.