

1.8. TRANSPORT PROPERTIES

The peak of the Lorentzian is at ε_0 and the width is γ . Fig. 1.8.5.1 shows the Seebeck coefficient calculated with this functional form. We arbitrarily took $\gamma = k_B T_\gamma$ and $T_\gamma = 100$ K. The four curves are for $\varepsilon_0/k_B = 10, 30, 100$ and 300 K, and the larger ε_0 has the larger Seebeck coefficient. All of the curves have a broad peak as a function of temperature. If $\varepsilon_0 = 0$, then $S = 0$, since the integrand is an odd function of $(\varepsilon - \mu)$. The vertical scale is S/S_0 where $S_0 = k_B/e = 86.17 \mu\text{V K}^{-1}$. We see from Fig. 1.8.5.1 that it is difficult to get values of S/S_0 very much larger than unity.

Another example is that of mixed-valence materials with f electrons. The f shells make electron states of narrow energy, which are approximated as Lorentzians. In this case, $\Sigma(\varepsilon)$ equals the inverse of the right-hand side of (1.8.5.10). The argument for this is that $1/\tau(\varepsilon)$ is proportional to the right-hand side of (1.8.5.10). Since $\Sigma \propto \tau$, it contains the inverse of (1.8.5.10). Interestingly enough, plots of the Seebeck coefficient by Jaccard & Sierro (1982) for this case also contain broad peaks in energy, where S/S_0 has a maximum of about unity. In this case, a proper calculation includes the fact that both C and γ are functions of temperature. For a review, see Mahan (1997).

We give these examples of the Seebeck coefficient because they are the cases that occur most often. In many metals, the Seebeck coefficient is either linear with temperature or has broad peaks. The broad peaks are due to structure in Σ near the chemical potential. This structure is usually due either to variations in the density of states or in the electron lifetime.

In insulators, the Seebeck coefficient can become relatively large. The exact value depends upon the energy gap, the temperature and the density of impurities. This example is treated in many references, e.g. Goldsmid (1986) and Rowe (1995).

1.8.6. Glossary

T	temperature (K)
\mathbf{J}	current density (A m^{-2})
\mathbf{J}_Q	heat current (W m^{-2})
σ	electrical conductivity (S m^{-1})
ρ	electrical resistivity
K	thermal conductivity ($\text{W m}^{-1} \text{K}^{-1}$)
\mathbf{E}	electric field (V m^{-1})
S	Seebeck coefficient (V K^{-1})
k_B	Boltzmann constant
R_H	Hall constant
m^*	effective mass of the electron
τ	lifetime of the electron

Research support from the University of Tennessee and from Oak Ridge National Laboratory, managed by Lockheed Martin Energy Research Corporation for the US Department of Energy under contract No. DE-AC05-96OR22464, is gratefully acknowledged.

References

- Allen, P. B., Beaulac, T. P., Khan, F. S., Butler, W. H., Pinski, F. J. & Swihart, J. C. (1986). *D.c. transport in metals*. *Phys. Rev. B*, **34**, 4331–4333.
- Allen, P. B., Berger, H., Chauvet, O., Forro, L., Jarlborg, T., Junod, A., Revaz, B. & Santi, G. (1996). *Transport properties, thermodynamic properties, and electronic structure of SrRuO₃*. *Phys. Rev. B*, **53**, 4393–4398.
- Allen, P. B. & Chakraborty, B. (1981). *Infrared and d.c. conductivity in metals with strong scattering: nonclassical behavior from a generalized Boltzmann equation containing band-mixing effects*. *Phys. Rev. B*, **23**, 4815–4827.
- Anthony, T. R., Banholzer, W. F., Fleischer, J. F., Wei, L., Kuo, P. K., Thomas, R. L. & Pryor, R. W. (1990). *Thermal diffusivity of isotropically enriched ¹²C diamond*. *Phys. Rev. B*, **42**, 1104–1111.
- Bass, J., Pratt, W. P. & Schroeder, P. A. (1990). *The temperature dependent electrical resistivities of the alkali metals*. *Rev. Mod. Phys.* **62**, 645–744.
- Berman, R. (1976). *Thermal conduction in solids*. Oxford University Press.
- Bruls, G. J. C. L., Bass, J., van Gelder, A. P., van Kempen, H. & Wyder, P. (1985). *Linear magnetoresistance due to sample thickness variations: applications to aluminum*. *Phys. Rev. B*, **32**, 1927–1939.
- Goldsmid, H. J. (1986). *Electronic refrigeration*. London: Pion Limited.
- Grimvall, G. (1981). *The electron–phonon interaction in metals*. Amsterdam: North-Holland.
- Jaccard, D. & Sierro, J. (1982). *Thermoelectric power of some intermediate valence compounds*. In *Valence instabilities*, edited by P. Wachter & H. Boppert, pp. 409–413. Amsterdam: North-Holland.
- Jacoboni, C., Canali, C., Ottaviani, G. & Quaranta, A. A. (1977). *A review of some charge transport properties of silicon*. *Solid State Electron.* **20**, 77–89.
- Jonson, M. & Girvin, S. M. (1979). *Electron–phonon dynamics and transport anomalies in random metal alloys*. *Phys. Rev. Lett.* **43**, 1447–1451.
- Klein, L., Dodge, J. S., Ahn, C. H., Snyder, G. J., Geballe, T. H., Beasley, M. R. & Kapitulnik, A. (1996). *Anomalous spin scattering effects in the badly metallic itinerant ferromagnet SrRuO₃*. *Phys. Rev. Lett.* **77**, 2774–2777.
- Klemens, P. G. (1955). *The scattering of low-frequency lattice waves by static imperfections*. *Proc. Phys. Soc. London Sect. A*, **68**, 1113–1128.
- Koshino, S. (1960). *Scattering of electrons by the thermal motion of impurity ions*. *Prog. Theor. Phys.* **24**, 484–494.
- MacDonald, A. H., Taylor, R. & Geldart, D. J. W. (1981). *Umklapp electron–electron scattering and the low-temperature electrical resistivity of the alkali metals*. *Phys. Rev. B*, **23**, 2718–2730.
- Mahan, G. D. (1990). *Many-particle physics*, 2nd ed. New York: Plenum.
- Mahan, G. D. (1997). *Good thermoelectrics*. In *Solid state physics*, Vol. 51, edited by H. Ehrenreich & F. Spaepen, pp. 81–157. New York: Academic Press.
- Mahan, G. D. & Roth, W. L. (1976). Editors. *Superionic conductors*. New York: Plenum.
- Mahan, G. D. & Wang, Z. (1989). *Koshino–Taylor coefficient in electrical conductivity*. *Phys. Rev. B*, **39**, 4926–4929.
- Matthiessen, A. & Vogt, C. (1864). *Ann. Phys. (Leipzig)*, **129**, 19.
- Mooij, J. H. (1973). *Electrical conduction in concentrated disordered transition metal alloys*. *Phys. Status Solidi A*, **17**, 521–530.
- Rode, D. L. (1972). *Electron mobility in Ge, Si, and GaP*. *Phys. Status Solidi B*, **53**, 245–254.
- Rode, D. L. (1975). *Low field electron transport*. In *Semiconductors and semimetals*, Vol. 10, edited by R. K. Willardson & A. C. Beer, pp. 1–89. New York: Academic Press.
- Rowe, D. M. (1995). Editor. *CRC handbook of thermoelectrics*. New York: CRC Press.
- Salamon, M. B. (1979). Editor. *Physics of superionic conductors*. New York: Springer-Verlag.
- Sievers, A. J. (1980). *Infrared and optical properties of Na, K, and Rb metals*. *Phys. Rev. B*, **22**, 1600–1611.
- Slack, G. A. (1979). *The thermal conductivity of nonmetallic crystals*. In *Solid state physics*, Vol. 34, edited by H. Ehrenreich, F. Seitz & D. Turnbull, pp. 1–71. New York: Academic Press.
- Spitzer, D. P. (1970). *Lattice thermal conductivity of semiconductors: a chemical bond approach*. *J. Phys. Chem. Solids*, **31**, 19–40.
- Taylor, P. L. (1964). *Changes in electrical resistance caused by incoherent electron–photon scattering*. *Phys. Rev. A*, **135**, 1333–1335.
- Tsuei, C. C. (1986). *Nonuniversality of the Mooij correlation – the temperature coefficient of electrical resistivity of disordered metals*. *Phys. Rev. Lett.* **57**, 1943–1946.
- Wiedemann, G. & Franz, R. (1853). *Ann. Phys. (Leipzig)*, **89**, 497.
- Wiser, N. (1984). *The electrical resistivity of simple metals*. *Contemp. Phys.* **25**, 211–249.
- Wu, J. W. & Mahan, G. D. (1984). *Transport equation at finite frequency: infrared absorption in metals*. *Phys. Rev. B*, **29**, 1769–1782.
- Yao, T. (1987). *Thermal properties of AlAs/GaAs superlattices*. *Appl. Phys. Lett.* **51**, 1798–1782.
- Ziman, J. M. (1962). *Electrons and phonons*. Oxford University Press.