

$\bar{6}c2$

$\bar{6}m2$



Hexagonal

No. 72

$\bar{6}c2$

Patterson symmetry $\bar{6}/mmm$

FIRST SETTING



Origin on $\bar{6}c1$

Asymmetric unit $0 \leq x; 0 \leq y; 0 \leq z \leq \frac{1}{4}$

Symmetry operations

- | | | |
|----------------------------------|----------------------------------|----------------------------------|
| (1) 1 | (2) $3^+ 0, 0, z$ | (3) $3^- 0, 0, z$ |
| (4) $m x, y, 0$ | (5) $\bar{6}^- 0, 0, z; 0, 0, 0$ | (6) $\bar{6}^+ 0, 0, z; 0, 0, 0$ |
| (7) $c x, \bar{x}, z$ | (8) $c x, 2x, z$ | (9) $c 2x, x, z$ |
| (10) $2 x, \bar{x}, \frac{1}{4}$ | (11) $2 x, 2x, \frac{1}{4}$ | (12) $2 2x, x, \frac{1}{4}$ |

Generators selected (1); $t(0,0,1)$; (2); (4); (7)

Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates			Reflection conditions		
				General:		
12 f 1	(1) x, y, z	(2) $\bar{y}, x - y, z$	(3) $\bar{x} + y, \bar{x}, z$	$l : l = 2n$		
	(4) x, y, \bar{z}	(5) $\bar{y}, x - y, \bar{z}$	(6) $\bar{x} + y, \bar{x}, \bar{z}$			
	(7) $\bar{y}, \bar{x}, z + \frac{1}{2}$	(8) $\bar{x} + y, y, z + \frac{1}{2}$	(9) $x, x - y, z + \frac{1}{2}$			
	(10) $\bar{y}, \bar{x}, \bar{z} + \frac{1}{2}$	(11) $\bar{x} + y, y, \bar{z} + \frac{1}{2}$	(12) $x, x - y, \bar{z} + \frac{1}{2}$			
				Special: no extra conditions		
6 e $m..$	$x, y, 0$	$\bar{y}, x - y, 0$	$\bar{x} + y, \bar{x}, 0$	$\bar{y}, \bar{x}, \frac{1}{2}$	$\bar{x} + y, y, \frac{1}{2}$	$x, x - y, \frac{1}{2}$
6 d $..2$	$x, \bar{x}, \frac{1}{4}$	$x, 2x, \frac{1}{4}$	$2\bar{x}, \bar{x}, \frac{1}{4}$	$x, \bar{x}, \frac{3}{4}$	$x, 2x, \frac{3}{4}$	$2\bar{x}, \bar{x}, \frac{3}{4}$
4 c $3..$	$0, 0, z$	$0, 0, \bar{z}$	$0, 0, z + \frac{1}{2}$	$0, 0, \bar{z} + \frac{1}{2}$		
2 b $\bar{6}..$	$0, 0, 0$	$0, 0, \frac{1}{2}$				
2 a 3.2	$0, 0, \frac{1}{4}$	$0, 0, \frac{3}{4}$				

Symmetry of special projections

Along [001] $3m$

Along [100] $\bar{6}1m1$

Along [210] $\bar{6}2mg$

$\mathbf{a}' = \frac{1}{2}\mathbf{c}$

$\mathbf{a}' = \mathbf{c}$

Origin at $0, 0, z$

Origin at $x, 0, 0$

Origin at $x, \frac{1}{2}x, \frac{1}{4}$

Maximal non-isotypic non-enantiomorphic subgroups

I	$[2]\bar{6}11 (\bar{6}, 59)$	1; 2; 3; 4; 5; 6
	$[2]\bar{6}3c1 (50)$	1; 2; 3; 7; 8; 9
	$[2]\bar{6}312 (46)$	1; 2; 3; 10; 11; 12
	$[3]\bar{6}mc2 (\bar{6}2cm, 19)$	1; 4; 7; 10
	$[3]\bar{6}mc2 (\bar{6}2cm, 19)$	1; 4; 8; 11
	$[3]\bar{6}mc2 (\bar{6}2cm, 19)$	1; 4; 9; 12

IIa none

IIb none

Maximal isotypic subgroups and enantiomorphic subgroups of lowest index

IIc $[3]\bar{6}c2 (\mathbf{c}' = 3\mathbf{c}) (72)$

Minimal non-isotypic non-enantiomorphic supergroups

I $[2]\bar{6}/mcc (74)$; $[2]\bar{6}_3/mmc (75)$

II $[2]\bar{6}m2 (\mathbf{c}' = \frac{1}{2}\mathbf{c}) (71)$



$\bar{6}2c$ $\bar{6}2m$

Hexagonal

No. 72

 $\bar{6}2c$ Patterson symmetry $\bar{6}/mmm$

SECOND SETTING

Origin on $\bar{6}1c$ Asymmetric unit $0 \leq x; 0 \leq y; 0 \leq z \leq \frac{1}{2}; y \leq x$

Symmetry operations

- | | | |
|-------------------------|--------------------------------|--------------------------------|
| (1) 1 | (2) 3^+ $0,0,z$ | (3) 3^- $0,0,z$ |
| (4) m $x,y,0$ | (5) $\bar{6}^-$ $0,0,z; 0,0,0$ | (6) $\bar{6}^+$ $0,0,z; 0,0,0$ |
| (7) 2 $x,x,\frac{1}{4}$ | (8) 2 $x,0,\frac{1}{4}$ | (9) 2 $0,y,\frac{1}{4}$ |
| (10) c x,x,z | (11) c $x,0,z$ | (12) c $0,y,z$ |

Generators selected (1); $t(0,0,1)$; (2); (4); (7)

Positions

Multiplicity, Wyckoff letter, Site symmetry	Coordinates			Reflection conditions
				General:
12 f 1	(1) x, y, z	(2) $\bar{y}, x - y, z$	(3) $\bar{x} + y, \bar{x}, z$	$l : l = 2n$
	(4) x, y, \bar{z}	(5) $\bar{y}, x - y, \bar{z}$	(6) $\bar{x} + y, \bar{x}, \bar{z}$	
	(7) $y, x, \bar{z} + \frac{1}{2}$	(8) $x - y, \bar{y}, \bar{z} + \frac{1}{2}$	(9) $\bar{x}, \bar{x} + y, \bar{z} + \frac{1}{2}$	
	(10) $y, x, z + \frac{1}{2}$	(11) $x - y, \bar{y}, z + \frac{1}{2}$	(12) $\bar{x}, \bar{x} + y, z + \frac{1}{2}$	
				Special: no extra conditions
6 e $m..$	$x, y, 0$	$\bar{y}, x - y, 0$	$\bar{x} + y, \bar{x}, 0$	$y, x, \frac{1}{2}$ $x - y, \bar{y}, \frac{1}{2}$ $\bar{x}, \bar{x} + y, \frac{1}{2}$
6 d $.2.$	$x, 0, \frac{1}{4}$	$0, x, \frac{1}{4}$	$\bar{x}, \bar{x}, \frac{1}{4}$	$x, 0, \frac{3}{4}$ $0, x, \frac{3}{4}$ $\bar{x}, \bar{x}, \frac{3}{4}$
4 c $3..$	$0, 0, z$	$0, 0, \bar{z} + \frac{1}{2}$	$0, 0, \bar{z}$	$0, 0, z + \frac{1}{2}$
2 b $\bar{6}..$	$0, 0, 0$	$0, 0, \frac{1}{2}$		
2 a $32.$	$0, 0, \frac{1}{4}$	$0, 0, \frac{3}{4}$		

Symmetry of special projections

Along [001] $3m$

Along [100] $\bar{6}2mg$

Along [210] $\bar{6}1m1$

$\mathbf{a}' = \mathbf{c}$

$\mathbf{a}' = \frac{1}{2}\mathbf{c}$

Origin at $0, 0, z$

Origin at $x, 0, \frac{1}{4}$

Origin at $x, \frac{1}{2}x, 0$

Maximal non-isotypic non-enantiomorphic subgroups

I	$[2]\bar{6}11 (\bar{6}, 59)$	1; 2; 3; 4; 5; 6
	$[2]\bar{6}31c (\bar{6}3c1, 50)$	1; 2; 3; 10; 11; 12
	$[2]\bar{6}321 (\bar{6}312, 46)$	1; 2; 3; 7; 8; 9
	$[3]\bar{6}m2c (\bar{6}2cm, 19)$	1; 4; 7; 10
	$[3]\bar{6}m2c (\bar{6}2cm, 19)$	1; 4; 8; 11
	$[3]\bar{6}m2c (\bar{6}2cm, 19)$	1; 4; 9; 12

IIa none

IIb none

Maximal isotypic subgroups and enantiomorphic subgroups of lowest index

IIc $[3]\bar{6}2c (\mathbf{c}' = 3\mathbf{c}) (\bar{6}c2, 72)$

Minimal non-isotypic non-enantiomorphic supergroups

I $[2]\bar{6}/mcc (74)$; $[2]\bar{6}_3/mmc (75)$

II $[2]\bar{6}m2 (\mathbf{c}' = \frac{1}{2}\mathbf{c}) (71)$