

Laue class  $C_{2h} - 2/m$ 

6. SCANNING TABLES

Monoclinic

**Laue class  $C_{2h} - 2/m$** 
**Geometric class  $C_2 - 112$** 

 No. 3  $P2$ 
 $C_2^1$ 

$$\mathcal{G} = P121 \quad \text{UNIQUE AXIS } b$$

$$\mathcal{G} = P112 \quad \text{UNIQUE AXIS } c$$

Orientation orbit ( <i>hkl</i> )	Conventional basis of the scanning group <b>a'</b> <b>b'</b> <b>d</b>	Scanning group $\mathcal{H}$	Linear orbit <b>sd</b>	Sectional layer group $\mathcal{L}(\mathbf{sd})$	
UNIQUE AXIS <i>b</i> (010) UNIQUE AXIS <i>c</i> (001)	<b>c</b> <b>a</b> <b>b</b> <b>a</b> <b>b</b> <b>c</b>	$P112$	<b>sd</b>	$p112$	L03
UNIQUE AXIS <i>b</i> ( <i>n0m</i> ) UNIQUE AXIS <i>c</i> ( <i>mn0</i> )	<b>b</b> $nc - ma$ $pc + qa$ <b>c</b> $na - mb$ $pa + qb$	$P211$	$0\mathbf{d}, \frac{1}{2}\mathbf{d}$ $[\mathbf{sd}, -\mathbf{sd}]$	$p211$ $p1$	L08 L01

 No. 4  $P2_1$ 
 $C_2^2$ 

$$\mathcal{G} = P12_11 \quad \text{UNIQUE AXIS } b$$

$$\mathcal{G} = P112_1 \quad \text{UNIQUE AXIS } c$$

Orientation orbit ( <i>hkl</i> )	Conventional basis of the scanning group <b>a'</b> <b>b'</b> <b>d</b>	Scanning group $\mathcal{H}$	Linear orbit <b>sd</b>	Sectional layer group $\mathcal{L}(\mathbf{sd})$	
UNIQUE AXIS <i>b</i> (010) UNIQUE AXIS <i>c</i> (001)	<b>c</b> <b>a</b> <b>b</b> <b>a</b> <b>b</b> <b>c</b>	$P112_1$	$[\mathbf{sd}, (s + \frac{1}{2})\mathbf{d}]$	$p1$	L01
UNIQUE AXIS <i>b</i> ( <i>n0m</i> ) UNIQUE AXIS <i>c</i> ( <i>mn0</i> )	<b>b</b> $nc - ma$ $pc + qa$ <b>c</b> $na - mb$ $pa + qb$	$P2_111$	$0\mathbf{d}, \frac{1}{2}\mathbf{d}$ $[\mathbf{sd}, -\mathbf{sd}]$	$p2_111$ $p1$	L09 L01