

4. CRYSTALLIZATION

glucopyranosides or alkyl- β -D-maltosides as detergents: the commercially available detergents are often 'contaminated' with the α -anomers in varying, sometimes substantial, concentrations. The α -anomers are much less soluble, and appear to prevent crystallization. In the case of photosystem I from a thermophilic cyanobacterium, it has been reported that for a 2% α -anomer content in dodecyl- β -D-maltoside preparations no crystals can be obtained, with a 0.5–2% content the diffraction of the crystals is anisotropic with a reduction in resolution to 5–6 Å, whereas diffraction to better than 3.5 Å resolution in all directions is observed when the content of the α -anomer is below 0.1% (Fromme & Witt, 1998). The percentage of α -anomer can be determined using NMR spectroscopy or high-performance liquid chromatography. During ageing of detergent solutions, a conversion from the β -anomer to the α -anomer is expected. Therefore, ageing of detergent solutions has to be prevented.

The detergents that have been successfully used to crystallize membrane proteins can also be found in Table 4.2.1.1. The possibilities for developing new detergents for membrane-protein crystallization have not been exhausted. There is a need for new detergents, e.g. detergents with head groups with sizes between glucose and maltose are still missing!

It has been observed that crystals of the photosynthetic reaction centre from the purple bacterium *Rhodospseudomonas viridis* could be obtained when *N,N*-dimethyldodecylamine-*N*-oxide is used as detergent, but not when *N,N*-dimethyldecylamine-*N*-oxide is used. Even crystals formed with the dodecyl homologue lost their order when soaked in a buffer containing the decyl homologue. These observations were found to have an obvious explanation when the location of the detergent molecules bound to the protein was determined using neutron crystallography (Roth *et al.*, 1989); the detergent molecules surrounding neighbouring photosynthetic reaction centres in the crystal lattice are in contact. It is likely that attractive interactions between neighbouring protein-bound detergent micelles contribute to the stability of the crystal lattice. Particularly striking (see Table 4.2.3.1) is the dependence of the crystal quality on the alkyl-chain length in the case of the two-subunit cytochrome *c* oxidase from the soil bacterium *Paracoccus denitrificans*. Well ordered crystals were obtained with undecyl- β -D-maltoside, but not with the decyl and dodecyl homologues. Table 4.2.3.1 also lists the names of important vendors of detergents.

4.2.4. The 'small amphiphile concept'

From the arguments and observations presented above, it is evident that the size and shape of the detergent micelle are very important in membrane-protein crystallization. Detergent micelles can be made smaller (and their curvatures changed) when small amphiphilic molecules like heptane-1,2,3-triol are added (Timmins *et al.*, 1991; Gast *et al.*, 1994). These compounds form mixed micelles with detergents. When 10% (*w/v*) heptane-1,2,3-triol is added to 1% solutions of *N,N*-dimethyldodecylamine-*N*-oxide in water, the number of detergent molecules per micelle decreases from 69 to 34, ~23 heptane-1,2,3-triol molecules are incorporated and the radius of the micelle is reduced from 20.9 to 16.9 Å (Timmins *et al.*, 1994). This so-called 'small amphiphile concept' has been used successfully to crystallize bacterial photosynthetic reaction centres (Michel, 1982, 1983; Buchanan *et al.*, 1993), bacterial light-harvesting complexes (Michel, 1991; Koepke *et al.*, 1996) and other membrane proteins (see Table 4.2.1.1). The light-harvesting complexes from the purple bacterium *Rhodospirillum rubrum* yield an astonishing number of different crystal forms, but only one diffracts to high resolution. A large amount of heptane-1,2,3-triol had to be added to obtain this crystal form. As a result, heptane-1,2,3-triol itself reached supersaturation during crystallization of

Table 4.2.3.1. Summary of the results of attempts to crystallize the two-subunit cytochrome *c* oxidase from the soil bacterium *Paracoccus denitrificans* using different detergents (after Ostermeier *et al.*, 1997)

The resolutions of the crystals obtained are given in parentheses. Abbreviations: C₁₂: dodecyl; C₁₁: undecyl; C₁₀: decyl; C₉: nonyl; CYMAL-6: (cyclohexyl)hexyl- β -D-maltoside; CYMAL-5: (cyclohexyl)pentyl- β -D-maltoside. *x* in E_{*x*} is the number of oxyethylene units in the alkylpolyoxyethylene detergents. Suppliers: A: Anatrace (Maumee, OH); B: Biomol; C: Calbiochem; F: Fluka.

Detergent	Supplier	Crystals?
C ₁₂ - β -D-maltoside	B	Yes (8 Å)
C ₁₁ - β -D-maltoside	B	Yes (2.5 Å)
C ₁₀ - β -D-maltoside	B	No
CYMAL-6	A	Yes (2.6 Å)
CYMAL-5	A	No
Dodecylsucrose	C	No
Decylsucrose	C	No
C ₉ - β -D-glucoside	C	No
C ₁₂ E ₈	F	Yes (> 8 Å)
C ₁₂ E ₆	F	No
C ₁₂ E ₅	F	No
C ₁₀ E ₆	F	No
C ₁₀ E ₅	F	No

the protein. Normally, the protein crystallizes first and heptane-1,2,3-triol later. When heptane-1,2,3-triol crystals form, the protein crystals crack and eventually redissolve. A likely explanation is that the concentration of solubilized heptane-1,2,3-triol is reduced when it forms crystals and the mixed detergent–heptane-1,2,3-triol micelles lose heptane-1,2,3-triol and take up detergent molecules from solution. The micelles expand and the crystals crack. This behaviour caused severe problems when searching for heavy-atom derivatives. Small amphiphiles work well with the *N,N*-dimethyl-alkylamine-*N*-oxides and alkylglucopyranosides as detergents, but not with alkylmaltosides. The most successful small amphiphiles are heptane-1,2,3-triol and benzamide.

4.2.5. Membrane-protein crystallization with the help of antibody Fv fragments

The number of detergents that can be tried is limited for rather unstable membrane proteins. For instance, the four-subunit cytochrome *c* oxidase from *P. denitrificans* is sufficiently stable only with dodecyl- β -D-maltoside as detergent. When other detergents are employed, subunits III and IV and some lipids are removed from subunits I and II. These lipids might contribute to the binding of the protein subunits III and IV to the central subunits I and II. Therefore, the size of the detergent micelles can not be varied by using different detergents. In order to obtain crystals, the size of the extramembraneous part of this important enzyme was enlarged by the binding of Fv fragments of monoclonal antibodies. For this purpose, monoclonal antibodies against the four-subunit cytochrome *c* oxidase were generated using the classical hybridoma technique. Then hybridoma cell lines producing conformation-specific antibodies were selected (such antibodies react positively in enzyme-linked immunosorbent assays, but negatively in Western blot assays). The cDNA strands coding for the respective VL and VH genes were cloned and expressed in *Escherichia coli*. Binding of conformation-specific antibody fragments can be expected to lead to a more homogeneous protein preparation. The first Fv fragment worked, and well ordered crystals of the four-subunit and