



14. ANOMALOUS DISPERSION


Figure 14.1.3.1

(a) Harker construction for a double isomorphous replacement. φ_M is the 'most probable' phase for \mathbf{F}_p . (b) Phase probability distribution corresponding to the double isomorphous replacement shown in part (a). The curve for derivative 1 is solid, that for derivative 2 is dashed, and that for the combined distribution is drawn as a dotted-and-dashed line.

From the heavy-atom parameters, the corresponding structure factor $\mathbf{F}_H(\mathbf{h})$ is calculated. To determine φ , the phase of $\mathbf{F}_p(\mathbf{h})$, we construct a set of phase circles, as proposed by Harker (1956). From a chosen origin O (Fig. 14.1.2.1a), the vector OA is drawn equal to $-\mathbf{F}_H$. Circles of radius F_p and F_{PH} are then drawn about O and A , respectively. The intersections of the phase circles at B and B' define two possible phase angles for F_p . Note that the angles are symmetrical about \mathbf{F}_H . This ambiguity may in principle be resolved in two ways: (a) by using a second heavy-atom isomorphous derivative or (b) by utilizing the anomalous-scattering effects for the first isomorph.

14.1.3. The method of multiple isomorphous replacement

The phase information provided by a second isomorph is illustrated in Fig. 14.1.3.1(a). In theory, the three phase circles will

intersect at a point and the phase ambiguity will be resolved. In practice, there will be errors in the observed amplitudes F_p and F_{PH} and in the heavy-atom parameters (and thus in \mathbf{F}_H). Also, the isomorphism may be imperfect. As a result, the intersections of the three phase circles may not coincide. Another complication arises from the fact that for reflections where \mathbf{F}_H is small, the circles will be essentially concentric and will not have well defined points of intersection. In other words, the phase determination will become indeterminate. The method of Blow & Crick (1959) was introduced as a way to take all these factors into account. It has had an extraordinary impact, not only as a practical method for protein phase determination, but also in influencing all subsequent thinking in this area.

14.1.4. The method of Blow & Crick


Blow & Crick pointed out that in practice the phase angle φ can never be determined with complete certainty. Rather, there is a finite probability that any arbitrary phase angle may be the correct one. Consider the vector diagram shown in Fig. 14.1.4.1, in which \mathbf{F}_H is known and we wish to determine the probability $P(\varphi)$ that the arbitrary phase angle φ is the correct phase of \mathbf{F}_p . Strictly, one should allow for the possibility of errors in \mathbf{F}_H , F_p and F_{PH} , and should consider the probability that the vector \mathbf{F}_p occupies all possible positions in the Argand diagram. However, Blow & Crick suggested that the analysis might be considerably simplified by assuming that F_p and \mathbf{F}_H are known accurately and that all the error lies in the observation of F_{PH} . In other words, it was assumed that the vector \mathbf{F}_p must lie on the circle of radius F_p , and the probability distribution of F_p could be evaluated as a function of φ only.

For an arbitrary phase angle φ , the phase triangle (Fig. 14.1.4.1) will not close exactly. If we define F_C to be the vector sum of \mathbf{F}_H and $F_p \exp(i\varphi)$, then the lack of closure of the phase triangle is given by

$$\varepsilon = F_C - F_{PH}. \quad (14.1.4.1)$$

Following Blow & Crick, if E is the r.m.s. error associated with the measurements, and the distribution of error is assumed to be Gaussian, then the probability $P(\varphi)$ of the phase φ being the true phase is

$$P(\varphi) = N \exp(-\varepsilon^2/2E^2), \quad (14.1.4.2)$$


Figure 14.1.4.1

Vector diagram illustrating the lack of closure, ε , of an isomorphous-replacement phase triangle.