

3.8. DATA CLUSTERING AND VISUALIZATION

Table 3.8.1

Six commonly used clustering methods

For each method, the coefficients α_i , β and γ in equation (3.8.11) are given.

Method	α_i	β	γ
Single link	$\frac{1}{2}$	0	$-\frac{1}{2}$
Complete link	$\frac{1}{2}$	0	$\frac{1}{2}$
Average link	$n_i/(n_i + n_j)$	0	0
Weighted-average link	$\frac{1}{2}$	0	0
Centroid	$n_i/(n_i + n_j)$	$-n_i n_j/(n_i + n_j)^2$	0
Sum of squares	$(n_i + n_k)/(n_i + n_j + n_k)$	$-n_k/(n_i + n_j + n_k)$	0

In some cases it can be advantageous to use $I^{1/2}$ in the distance-matrix generation; this can enhance the sensitivity of the clustering to weak peaks (Butler *et al.*, 2019).

3.8.3. Cluster analysis

Cluster analysis uses **d** (or **s**, or **δ**) to partition the patterns into groups based on the similarity of their diffraction profiles. Associated with cluster are a number of important ancillary techniques all of which will be discussed here. A flowchart of these methods is shown in Fig. 3.8.4.


3.8.3.1. Dendrograms

Using **d** and **s**, agglomerative, hierarchical cluster analysis is now carried out, in which the patterns are put into clusters as defined by their distances from each other. [Gordon (1981, 1999) and Everitt *et al.* (2001) provide excellent and detailed introductions to the subject. Note that the two editions of Gordon's monograph are quite distinct and complementary.] The method begins with a situation in which each pattern is considered to be in a separate cluster. It then searches for the two patterns with the shortest distance between them, and joins them into a single cluster. This continues in a stepwise fashion until all the patterns form a single cluster. When two clusters (C_i and C_j) are merged, there is the problem of defining the distance between the newly formed cluster $C_i \cup C_j$ and any other cluster C_k . There are a number of different ways of doing this, and each one gives rise to a different clustering of the patterns, although often the difference can be quite small. A general algorithm has been proposed by Lance & Williams (1967), and is summarized in a simplified form by Gordon (1981). The distance from the new cluster formed by merging C_i and C_j to any other cluster C_k is given by

$$d(C_i \cup C_j, C_k) = \alpha_i d(C_i, C_k) + \alpha_j d(C_j, C_k) + \beta d(C_i, C_j) + \gamma |d(C_i, C_k) - d(C_j, C_k)|. \quad (3.8.11)$$

There are many possible clustering methods. Table 3.8.1 defines six commonly used clustering methods, defined in terms of the parameters α , β and γ . All these methods can be used with powder data; in general, the group-average-link or single-link formalism is the most effective, although differences between the methods are often slight.

The results of cluster analysis are usually displayed as a dendrogram, a typical example of which is shown in Fig. 3.8.6(a), where a set of 13 powder patterns is analysed using the centroid method. Each pattern begins at the bottom of the plot as a separate cluster, and these amalgamate in stepwise fashion linked by horizontal tie bars. The height of the tie bar represents a similarity measure as measured by the relevant distance. As an

**Figure 3.8.2**

Four different methods of estimating the number of clusters present in a set of 23 powder patterns for the drug doxazosin. A total of five polymorphs are present, as well as two mixtures of these polymorphs. (a) A scree plot from the eigenvalue analysis of the correlation matrix; (b) the use of the C test (the coefficients have been multiplied by 100.0), which gives an estimate of five clusters using its local maximum. The γ test estimates that there are seven clusters and the CH test has a local maximum at seven clusters. Numerical details are given in Table 3.8.2.

indication of the differences that can be expected in the various algorithms used for dendrogram generation, Fig. 3.8.6(e) shows the same data analysed using the single-link method: the resulting clustering is slightly different: the similarity measures are larger, and, in consequence, the tie bars are higher on the graph. [For further examples see Barr *et al.* (2004b,c) and Barr, Dong, Gilmore & Faber (2004).]

3.8.3.2. Estimating the number of clusters

An estimate of the number of clusters present in the data set is needed. In terms of the dendrogram, this is equivalent to 'cutting the dendrogram' *i.e.* the placement of a horizontal line across it such that all the clusters as defined by tie lines above this line remain independent and unlinked. The estimation of the number of clusters is an unsolved problem in classification methods. It is easy to see why: the problem depends on how similar the patterns need to be in order to be classed as the same, and how much variability is allowed within a cluster. We use two approaches: (a) eigenvalue analysis of matrices **ρ** and **\mathbf{A}** , and (b) those based on cluster analysis.

Eigenvalue analysis is a well used technique: the eigenvalues of the relevant matrix are sorted in descending order and when a fixed percentage (typically 95%) of the data variability has been accounted for, the number of eigenvalues is selected. This is shown graphically *via* a scree plot, an example of which is shown in Fig. 3.8.2.

3. METHODOLOGY

We carry out eigenvalue analysis on the following:

- (1) Matrix ρ .
- (2) Matrix \mathbf{A} , as described in Section 3.8.3.3.
- (3) A transformed form of ρ in which ρ is standardized to give ρ_s in which the rows and columns have zero mean and unit variance. The matrix $\rho_s \rho_s^T$ is then computed and subjected to eigenanalysis. It tends to give a lower estimate of cluster numbers than (1).

The most detailed study on cluster counting is that of Milligan & Cooper (1985), and is summarized by Gordon (1999). From this we have selected three tests that seem to operate effectively with powder data:

- (4) The Calinški & Harabasz (1974) (CH) test:

$$CH(c) = [B/(c-1)]/[W/(n-c)]. \quad (3.8.12)$$

A centroid is defined for each cluster. W denotes the total within-cluster sum of squared distances about the cluster centroids, and B is the total between-cluster sum of squared distances. Parameter c is the number of clusters chosen to maximize CH.

- (5) A variant of Goodman & Kruskal's (1954) γ test, as described by Gordon (1999). The dissimilarity matrix is used. A comparison is made between all the within-cluster dissimilarities and all the between-cluster dissimilarities. Such a comparison is marked as concordant if the within-cluster dissimilarity is less than the between-cluster dissimilarity, and discrepant otherwise. Equalities, which are unusual, are disregarded. If S_+ is the number of concordant and S_- the number of discrepant comparisons, then

$$\gamma(c) = (S_+ - S_-)/(S_+ + S_-). \quad (3.8.13)$$

A maximum in γ is sought by an appropriate choice of cluster numbers.

- (6) The C test (Milligan & Cooper, 1985). This chooses the value of c that minimizes

$$C(c) = [W(c) - W_{\min}]/(W_{\max} - W_{\min}). \quad (3.8.14)$$

$W(c)$ is the sum of all the within-cluster dissimilarities. If the partition has a total of r such dissimilarities, then W_{\min} is the sum of the r smallest dissimilarities and W_{\max} is the sum of the r largest.

The results of tests (4)–(6) depend on the clustering method being used. To reduce the bias towards a given dendrogram method, these tests are carried out on four different clustering methods: the single-link, the group-average, the sum-of-squares and the complete-link methods. Thus there are 12 semi-independent estimates of the number of clusters from clustering methods, and three from eigenanalysis, making 15 in all.

A composite algorithm is used to combine these estimates. The maximum and minimum values of the number of clusters (c_{\max} and c_{\min} , respectively) given by the eigenanalysis results [(1)–(3) above] define the primary search range; tests (4)–(6) are then used in the range $\min(c_{\max} + 3, n) \leq c \leq \max(c_{\min} - 3, 0)$ to find local maxima or minima as appropriate. The results are averaged, any outliers are removed, and a weighted mean value is taken of the remaining indicators, then this is used as the final estimate of the number of clusters. Confidence levels for c are also defined by the estimates of the maximum and minimum cluster numbers after any outliers have been removed.

A typical set of results for the PXRD data from 23 powder patterns for doxazosin (an anti-hypertension drug) in which five polymorphs are present, as well as two mixtures of polymorphs, is shown in Fig. 3.8.2(a) and (b) (see also Table 3.8.2). The scree

Table 3.8.2

Estimate of the number of clusters for the 23 sample data set for doxazosin

There are five polymorphs present, plus two mixtures of these polymorphs. The maximum estimate is 7; the minimum estimate is 4; the combined weighted estimate of the number of clusters is 6, and the median value is 5. The dendrogram cut level is set to give 5 clusters, and the lower and upper confidence limits are 4 and 7, respectively.

Method	No. of clusters
Principal-component analysis (non-transformed matrix)	5
Principal-component analysis (transformed matrix)	4
Multidimensional metric scaling	4
γ statistic using single linkage	7
CH statistic using single linkage	7
C statistic using single linkage	—
γ statistic using group averages	7
CH statistic using group averages	5
C statistic using group averages	—
γ statistic using sum of squares	—
CH statistic using sum of squares	5
C statistic using sum of squares	—
γ statistic using complete linkage	—
CH statistic using complete linkage	5
C statistic using complete linkage	—

plot arising from the eigenanalysis of the correlation matrix indicates that 95% of the variability can be accounted for by five components, and this is shown in Fig. 3.8.2(a). Eigenvalues from other matrices indicate that four clusters are appropriate. A search for local optima in the CH, γ and C tests is then initiated in the range 2–8 possible clusters. Four different clustering methods are tried, and the results indicate a range of 4–7 clusters. There are no outliers, and the final weighted mean value of 5 is calculated. As Fig. 3.8.2(b) shows, the optimum points for the C and γ tests are often quite weakly defined (Barr *et al.*, 2004b).

3.8.3.3. Metric multidimensional scaling

This is, in its essentials, the particle-in-a-box problem. Each powder pattern is represented as a single sphere, and these spheres are placed in a cubic box of unit dimensions such that the positions of the spheres reproduce as closely as possible the distance matrix, \mathbf{d} , generated from correlating the patterns. The spheres have an arbitrary orientation in the box.

To do this, the $(n \times n)$ distance matrix \mathbf{d} is used in conjunction with metric multidimensional scaling (MMDS) to define a set of p underlying dimensions that yield a Euclidean distance matrix, \mathbf{d}^{calc} , whose elements are equivalent to or closely approximate the elements of \mathbf{d} .

The method works as follows (Cox & Cox, 2000; Gower, 1966; Gower & Dijksterhuis, 2004).

The matrix \mathbf{d} has zero diagonal elements, and so is not positive semidefinite. A positive definite matrix, $\mathbf{A}(n \times n)$ can be constructed, however, by computing

$$\mathbf{A} = -\frac{1}{2} \left[\mathbf{I}_n - \frac{1}{n} \mathbf{i}_n \mathbf{i}_n' \right] \mathbf{D} \left[\mathbf{I}_n - \frac{1}{n} \mathbf{i}_n \mathbf{i}_n' \right], \quad (3.8.15)$$

where \mathbf{I}_n is an $(n \times n)$ identity matrix, \mathbf{i}_n is an $(n \times 1)$ vector of unities and \mathbf{D} is defined in equation (3.8.8). The matrix $[\mathbf{I}_n - (1/n)\mathbf{i}_n \mathbf{i}_n']$ is called a centring matrix, since \mathbf{A} has been derived from \mathbf{D} by centring the rows and columns.

The eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ and the corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ are then obtained. A total of p eigenvalues of \mathbf{A} are positive and the remaining $(n - p)$ will be zero. For the p